arXiv:1501.03319v2 [math.DS] 23 Jan 2015

Random Iteration of Maps on a Cylinder and diffusive behavior

O. Castejón[*] V. Kaloshin[†]

January 26, 2015

Abstract

In this paper we propose a model of random compositions of maps of a cylinder, which in the simplified form is as follows: $(\theta, r) \in \mathbb{T}\times\mathbb{R} = \mathbb{A}$ and

$$f_{\pm 1} : \begin{pmatrix} \theta \\ r \end{pmatrix} \longmapsto \begin{pmatrix} \theta + r + \varepsilon u_{\pm 1}(\theta, r). \\ r + \varepsilon v_{\pm 1}(\theta, r). \end{pmatrix},$$

where $u_\pm$ and $v_\pm$ are smooth and $v_\pm$ are trigonometric polynomials in θ such that $\int v_\pm(\theta, r)\, d\theta = 0$ for each r. We study the random compositions

$$(\theta_n, r_n) = f_{\omega_{n-1}} \circ \cdots \circ f_{\omega_0}(\theta_0, r_0)$$

with $\omega_k \in \{-1, 1\}$ with equal probabilities. We show that under natural non-degeneracy hypothesis for $n \sim \varepsilon^{-2}$ the distributions of $r_n - r_0$ weakly converge to a diffusion process with explicitly computable drift and variance.

In the case of random iteration of the standard maps

$$f_{\pm 1} : \begin{pmatrix} \theta \\ r \end{pmatrix} \longmapsto \begin{pmatrix} \theta + r + \varepsilon v_{\pm 1}(\theta). \\ r + \varepsilon v_{\pm 1}(\theta) \end{pmatrix},$$

where $v_\pm$ are trigonometric polynomials such that $\int v_\pm(\theta)\, d\theta = 0$ we prove a vertical central limit theorem. Namely, for $n \sim \varepsilon^{-2}$ the distributions of $r_n - r_0$ weakly converge to a normal distribution $\mathcal{N}(0, \sigma^2)$ for $\sigma^2 = \frac{1}{4}\int (v_+(\theta) - v_-(\theta))^2\, d\theta$.

Such random models arise as a restrictions to a Normally Hyperbolic Invariant Lamination for a Hamiltonian flow of the generalized example of Arnold. We hope that this mechanism of stochasticity sheds some light on formation of diffusive behaviour at resonances of nearly integrable Hamiltonian systems.

---

[*] Universitat Politècnica de Catalunya, oriol.castejon@gmail.com

[†] University of Maryland at College Park, vadim.kaloshin@gmail.com

Contents

1 Introduction **3**
- 1.1 Motivation: Arnold diffusion and instabilities 3
- 1.2 KAM stability 3
- 1.3 A priori unstable systems 4
- 1.4 Random fluctuations of eccentricity in Kirkwood gaps in the asteroid belt 6
- 1.5 Random iteration of cylinder maps 6
- 1.6 Diffusion processes and infinitesimal generators 7

2 The model and statement of the main result **8**

3 Strategy of the proof **11**
- 3.1 Strip decomposition 11
- 3.2 Subdivision of the cylinder into domains with different quantitative behaviour 12
- 3.3 The normal form 13
- 3.4 Analsys of the Martingale problem in each kind of strip 14
  - 3.4.1 A Totally Irrational Strip 15
  - 3.4.2 An Imaginary Rational Strip 15
  - 3.4.3 A Real Rational Strip 16
  - 3.4.4 A transition zone 16
- 3.5 From the local diffusion in the rational strips to the global diffusion on the line 17
- 3.6 Plan of the rest of the paper 19

4 The Normal Form Theorem **19**

5 Analysis of the Martingale problem in the strips of each type **28**
- 5.1 The Totally Irrational case 28
- 5.2 The Imaginary Rational case 38
- 5.3 The Real Rational case 43
- 5.4 Transition Zones 50

A Measure of the domain covered by RR and IR intervals **54**

B Sufficient condition for weak convergence and auxiliary lemmas **55**

1 Introduction

1.1 Motivation: Arnold diffusion and instabilities

By Arnold-Louiville theorem a completely integrable Hamiltonian system can be written in action-angle coordinates, namely, for action p in an open set $U \subset \mathbb{R}^n$ and angle θ on an n-dimensional torus $\mathbb{T}^n$ there is a function $H_0(p)$ such that equations of motion have the form

$$\dot{\theta} = \omega(p), \quad \dot{p} = 0, \qquad \text{where } \ \omega(p) := \partial_p H_0(p).$$

The phase space is foliated by invariant n-dimensional tori $\{p = p_0\}$ with either periodic or quasi-periodic motions $\theta(t) = \theta_0 + t\,\omega(p_0)$ (mod 1). There are many different examples of integrable systems (see e.g. wikipedia).

It is natural to consider small Hamiltonian perturbations

$$H_\varepsilon(\theta, p) = H_0(p) + \varepsilon H_1(\theta, p), \qquad \theta \in \mathbb{T}^n, \ p \in U$$

where ε is small. The new equations of motion become

$$\dot{\theta} = \omega(p) + \varepsilon \partial_p H_1, \quad \dot{p} = -\varepsilon \partial_\theta H_1.$$

In the sixties, Arnold [1] (see also [2, 3]) conjectured that *for a generic analytic perturbation there are orbits $(\theta, p)(t)$ for which the variation of the actions is of order one, i.e. $\|p(t) - p(0)\|$ that is bounded from below independently of ε for all ε sufficiently small.*

See [5, 10, 25, 26, 27] about recent progress proving this conjecture for convex Hamiltonians.

1.2 KAM stability

Obstructions to Arnold diffusion, and to any form of instability in general, are widely known, following the works of Kolmogorov, Arnold, and Moser called nowadays KAM theory. The fundamental result says that for a properly nondegenerate H_0 and for all sufficiently regular perturbations εH_1, the system defined by H_ε still has many invariant n-dimensional tori. These tori are small deformation of unperturbed tori and measure of the union of these invariant tori tends to the full measure as ε goes to zero.

One consequence of KAM theory is that for $n = 2$ there are no instabilities. Indeed, generic energy surfaces $S_E = \{H_\varepsilon = E\}$ are 3-dimensional manifolds, KAM tori are 2-dimensional. Thus, KAM tori separate surfaces S_E and prevent orbits from diffusing.

1.3 A priori unstable systems

As an interesting model [1] Arnold proposed to study the following example

$$H_\varepsilon(p,q,I,\varphi,t) = \frac{I^2}{2} + H_0(p,q) + \varepsilon H_1(p,q,I,\varphi,t) :=$$
$$= \underbrace{\frac{I^2}{2}}_{harmonic\ oscillator} + \underbrace{\frac{p^2}{2} + (\cos q - 1)}_{pendulum} + \varepsilon H_1(p,q,I,\varphi,t), \tag{1}$$

where $q, \varphi, t \in \mathbb{T}$ are angles, $p, I \in \mathbb{R}$ are actions (see Fig. 3) and $H_1 = (\cos q - 1)(\cos\varphi + \cos t)$.



Figure 1: The rotor times the pendulum

For $\varepsilon = 0$ the system is a direct product of the harmonic oscillator $\ddot\varphi = 0$ and the pendulum $\ddot q = \sin q$. Instabilities occur when the (p,q)-component follows the separatrices $H_0(p,q) = 0$ and passes near the saddle $(p,q) = (0,0)$. Equations of motion for H_ε have a (normally hyperbolic) invariant cylinder Λ_ε which is $\mathcal{C}^1$ close to $\Lambda_0 = \{p = q = 0\}$. Systems having an invariant cylinder with a family of separatrix loops are called *an apriori unstable*. Since they were introduced by Arnold [1], they received a lot of attention both in mathematics and physics community see e.g. [4, 9, 10, 11, 13, 21, 39, 40].

Chirikov [10] and his followers made extensive numerical studies for the Arnold example. It indicates that *the I-displacement behaves randomly, where randomness is due to choice of initial conditions near* $H_0(p,q) = 0$.

More exactly, integration of solutions whose "initial conditions" randomly chosen ε-close to $H_0(p,q) = 0$ and integrated over time $\sim -\varepsilon^{-2}\ln\varepsilon$-time. This leads to the I- displacement being of order of one and having some distribution. This coined the name for this phenomenon: *Arnold diffusion.*

Let $\varepsilon = 0.01$ and $T = -\varepsilon^{-2}\ln\varepsilon$. On Fig. 1.3 we present several histograms plotting displacement of the I-component after time $T, 2T, 4T, 8T$ with 6 different groups of initial conditions, and histograms of 10^6 points. In each group we start with a large set of initial conditions close to $p = q = 0,\ I = I^*$.[1] One of the distinct features is that only one distribution (a) is close symmetric, while in all others have a drift.

---

[1] These histograms are part of the forthcoming paper of the second author with P. Roldan with extensive numerical analysis of dynamics of the Arnold's example.



(a)



(b)



(c)



(d)



(e)



(f)

A similar stochastic behaviour was observed numerically in many other nearly integrable problems ([10] pg. 370, [16, 28], see also [36]). To give another illustrative example consider motion of asteroids in the asteroid belt.

1.4 Random fluctuations of eccentricity in Kirkwood gaps in the asteroid belt

The asteroid belt is located between orbits of Mars and Jupiter and has around one million asteroids of diameter of at least one kilometer. When astronoters build a histogram based on orbital period of asteroids there are well known gaps in distribution called *Kirkwood gaps* (see Figure below).



These gaps occur when ratio of of an asteroid and of Jupiter is a rational with small denominator: $1/3, 2/5, 3/7$. This correspond to so called *mean motion resonances for the three body problem.* Wisdom [41] made a numerical analysis of dynamics at mean motion resonance and observed *random fluctuations of eccentricity* of asteroids. As these fluctuations grow and eccentricity reaches a certain critical value an orbit of a hypothetic asteroid starts to cross the orbit of Mars. This eventually leads either to a collision of the asteroid with Mars or a close encounter. The latter changes the orbit so drastically that almost certainly it disappears from the asteroid belt. In [17] we only managed to prove existence of certain orbits whose eccentricity change by $O(1)$ for the restricted planar three body problem. Outside of these resonances one could argue that KAM theory provides stability [32].

1.5 Random iteration of cylinder maps

Consider the time one map of H_ε, denoted

$$F_\varepsilon : (p, q, I, \varphi) \to (p', q', I', \varphi').$$

It turns out that for initial conditions ε-close to $H_0(p, q) = 0$, except of a hypersurface, one can define a return map to an $O(\varepsilon)$-neighborhood of $(p, q) = 0$. Often such a map is called *a separatrix map* and in the 2-dimensional case was introduced by physicists Filonenko-Zaslavskii [18]. In multidimensional setting such a map was defined and studied by Treschev [33, 38, 39, 40].

It turns starting near $(p,q)=0$ and iterating F_ε until the orbit comes back $(p,q)=0$ leads to a family of maps of a cylinder

$$f_{\varepsilon,p,q} : (I,\varphi) \to (I',\varphi'), \qquad (I,\varphi) \in \mathbb{A} = \mathbb{R} \times \mathbb{T}$$

which are close to integrable. Since at $(p,q)=0$ the (p,q)-component has a saddle, there is a sensitive dependence on initial condition in (p,q) and returns do have some randomness in (p,q). The precise nature of this randomness at the moment is not clear. There are several coexisting behaviours, including unstable diffusive, stable quasi-periodic, orbits can stick to KAM tori, and which one is dominant is to be understood. May be mechanism of capture into resonances [15] is also relevant in this setting.

In [22] we construct a normally hyperbolic lamination (NHL) for an open class of trigonometric perturbations of the form

$$H_1 = (\cos q - 1)P(\exp(i\varphi), \exp(it)).$$

Constructing unstable orbits along NHL is also discussed in [14]. In general, NHL give rise to a skew shift. For example, let $\Sigma = \{-1,1\}^{\mathbb{Z}}$ be the space of infinite sequences of -1's and 1's and $\sigma : \Sigma \to \Sigma$ be the standard shift.

Consider a skew product of cylinder maps

$$F : \mathbb{A} \times \Sigma \to \mathbb{A} \times \Sigma, \qquad F(r,\theta;\omega) = (f_\omega(r,\theta), \sigma\omega),$$

where each $f_\omega(r,\theta)$ *is a nearly integrable cylinder maps, in the sense that it almost preserves the* r*-component* [2]*.*

The goal of the present paper is to study a wide enough class of skew products so that they arise in Arnold's example with a trigonometric perturbation of the above type (see [22]).

Now we formalize our model and present the main result.

1.6 Diffusion processes and infinitesimal generators

In order to formalise the statement about diffusive behaviour we need to recall some basic probabilistic notions. Consider a Brownian motion $\{B_t,\, t \geq 0\}$.

A Brownian motion is a properly chosen limit of the standard random walk. A generalisation of a Brownian motion is *a diffusion process* or *an Ito diffusion.* To define it let (Ω, Σ, P) be a probability space. Let R :

---

[2] The reason we switch from the (I,φ)-coordinates on the cylinder to (r,θ) is because we perform a coordinate change.

$[0,+\infty) \times \Omega \to \mathbb{R}$. It is called an Ito diffusion if it satisfies *a stochastic differential equation* of the form

$$\mathrm{d}R_t = b(R_t)\,\mathrm{d}t + \sigma(R_t)\,\mathrm{d}B_t, \tag{2}$$

where B is an Brownian motion, $b : \mathbb{R} \to \mathbb{R}$ and $\sigma : \mathbb{R} \to \mathbb{R}$ are Lipschitz functions called the drift and the variance respectively. For a point $r \in \mathbb{R}$, let $\mathbb{P}_r$ denote the law of X given initial data $R_0 = r$, and let $\mathbb{E}_r$ denote expectation with respect to $\mathbb{P}_r$.

The *infinitesimal generator* of R is the operator A, which is defined to act on suitable functions $f : \mathbb{R} \to \mathbb{R}$ by

$$Af(r) = \lim_{t \downarrow 0} \frac{\mathbb{E}_r[f(R_t)] - f(r)}{t}.$$

The set of all functions f for which this limit exists at a point r is denoted $D_A(r)$, while D_A denotes the set of all f's for which the limit exists for all $r \in \mathbb{R}$. One can show that any compactly-supported $\mathcal{C}^2$ function f lies in D_A and that

$$Af(r) = b(r)\frac{\partial f}{\partial r} + \frac{1}{2}\sigma^2(r)\frac{\partial^2 f}{\partial r \partial r}. \tag{3}$$

The distribution of a diffusion process is characterise by the drift $b(r)$ and the variance $\sigma^2(r)$.

2 The model and statement of the main result

Let $\varepsilon > 0$ be a small parameter and $l \geq 12$ be an integer. Denote by $\mathcal{O}_l(\varepsilon)$ a $\mathcal{C}^l$ function whose $\mathcal{C}^l$ norm is bounded by $C\varepsilon$ with C independent of ε. Similar definition applies for a power of ε. As before Σ denotes $\{0,1\}^{\mathbb{Z}}$ and $\omega = (\dots, \omega_0, \dots) \in \Sigma$.

Consider two nearly integrable maps:

$$\begin{aligned} f_\omega : \mathbb{T} \times \mathbb{R} &\longrightarrow \mathbb{T} \times \mathbb{R} \\ f_\omega : \begin{pmatrix} \theta \\ r \end{pmatrix} &\longmapsto \begin{pmatrix} \theta + r + \varepsilon u_{\omega_0}(\theta, r) + \mathcal{O}_l(\varepsilon^{1+a}, \omega) \\ r + \varepsilon v_{\omega_0}(\theta, r) + \varepsilon^2 w_{\omega_0}(\theta, r) + \mathcal{O}_l(\varepsilon^{2+a}, \omega) \end{pmatrix}. \end{aligned} \tag{4}$$

for $\omega_0 \in \{-1, 1\}$, where u_{ω_0}, v_{ω_0}, and w_{ω_0} are bounded $\mathcal{C}^l$ functions, 1-periodic in θ, $\mathcal{O}_l(\varepsilon^{1+a}, \omega)$ and $\mathcal{O}_l(\varepsilon^{2+a}, \omega)$ denote remainders depending on ω and uniformly C^l bounded in ω, and $0 < a \leq 1/6$. Assume

$$\max |v_i(\theta, r)| \leq 1,$$

where maximum is taken over $i = -1, 1$ and all $(\theta, r) \in \mathbb{A}$, otherwise, renormalize ε.

We study random iterations of the maps f_1 and f_{-1}, such that at each step the probability of performing either map is $1/2$. Importance of understanding iterations of several maps for problems of diffusion is well known (see e.g. [24, 32]).

Denote the expected potential and the difference of potentials by

$$\mathbb{E}u(\theta, r) := \frac{1}{2}(u_1(\theta, r) + u_{-1}(\theta, r)), \quad \mathbb{E}v(\theta, r) := \frac{1}{2}(v_1(\theta, r) + v_{-1}(\theta, r)),$$

$$u(\theta, r) := \frac{1}{2}(u_1(\theta, r) - u_{-1}(\theta, r)), \quad v(\theta, r) := \frac{1}{2}(v_1(\theta, r) - v_{-1}(\theta, r)).$$

Suppose the following assumptions hold:

[H0] (*zero average*) Let for each $r \in \mathbb{R}$ and $i = \pm 1$ we have $\int v_i(\theta, r)\, d\theta = 0$.

[H1] (*no common zeroes*) For each integer $n \in \mathbb{Z}$ potentials $v_1(\theta, n)$ and $v_{-1}(\theta, n)$ have no common zeroes and, equivalently, f_1 and f_{-1} have no fixed points;

[H2] for each $r \in \mathbb{R}$ we have $\int_0^1 v^2(\theta, r) d\theta =: \sigma(r) \neq 0$;

[H3] The functions $v_i(\theta, r)$ are trigonometric polynomials in θ, i.e. for some positive integer d we have

$$v_i(\theta, r) = \sum_{k \in \mathbb{Z},\ |k| \leq d} v^{(k)}(r) \exp 2\pi i k \theta.$$

For $\omega \in \{-1, 1\}^{\mathbb{Z}}$ we can rewrite the maps f_ω in the following form:

$$f_\omega \begin{pmatrix} \theta \\ r \end{pmatrix} \longmapsto \begin{pmatrix} \theta + r + \varepsilon \mathbb{E}u(\theta, r) + \varepsilon \omega_0 u(\theta, r) + \mathcal{O}_l(\varepsilon^{1+a}, \omega) \\ r + \varepsilon \mathbb{E}v(\theta, r) + \varepsilon \omega_0 v(\theta, r) + \varepsilon^2 w_{\omega_0}(\theta, r) + \mathcal{O}_l(\varepsilon^{2+a}, \omega) \end{pmatrix}.$$

Let n be positive integer and $\omega_k \in \{-1, 1\}$, $k = 0, \ldots, n-1$, be independent random variables with $\mathbb{P}\{\omega_k = \pm 1\} = 1/2$ and $\Omega_n = \{\omega_0, \ldots, \omega_{n-1}\}$. Given an initial condition (θ_0, r_0) we denote:

$$(\theta_n, r_n) := f^n_{\Omega_n}(\theta_0, r_0) = f_{\omega_{n-1}} \circ f_{\omega_{n-2}} \circ \cdots \circ f_{\omega_0}(\theta_0, r_0). \tag{5}$$

[H4] (*no common periodic orbits*) Suppose for any rational $r = p/q \in \mathbb{Q}$ with p, q relatively prime, $1 \leq |q| \leq 2d$ and any $\theta \in \mathbb{T}$

$$\sum_{k=1}^{q} \left[v_{-1}(\theta + \frac{k}{q}, r) - v_1(\theta + \frac{k}{q}, r) \right]^2 \neq 0.$$

This prohibits f_1 and f_{-1} to have common periodic orbits of period $|q|$.

[H5] (*no degenerate periodic points*) Suppose for any rational $r = p/q \in \mathbb{Q}$ with p, q relatively prime, $1 \leq |q| \leq d$, the function:

$$\mathbb{E}v_{p,q}(\theta, r) = \sum_{\substack{k \in \mathbb{Z} \\ 0 < |kq| < d}} \mathbb{E}v^{kq}(r) e^{2\pi i k q \theta}$$

has distinct non-degenerate zeroes, where $\mathbb{E}v^j(r)$ denotes the j–th Fourier coefficient of $\mathbb{E}v(\theta, r)$.

A straightforward calculation shows that:

$$\begin{aligned}
\theta_n &= \theta_0 + n r_0 + \varepsilon \sum_{k=0}^{n-1} \left(\mathbb{E}u(\theta_k, r_k) + \mathbb{E}v(\theta_k, r_k) \right) \\
&\quad + \varepsilon \sum_{k=0}^{n-1} \omega_k \left(u(\theta_k, r_k) + v(\theta_k, r_k) \right) + \mathcal{O}_l(n\varepsilon^{1+a}) \qquad (6) \\
r_n &= r_0 + \varepsilon \sum_{k=0}^{n-1} \mathbb{E}v(\theta_k, r_k) + \varepsilon \sum_{k=0}^{n-1} \omega_k v(\theta_k, r_k) + \mathcal{O}_l(n\varepsilon^{2+a})
\end{aligned}$$

Even though these maps might not be area-preserving, using normal forms we will simplify these maps significantly on a large domain of the cylinder.

Theorem 2.1. *Assume that in the notations above conditions* **[H0-H5]** *hold. Let $n_\varepsilon \varepsilon^2 \to s > 0$ as $\varepsilon \to 0$ for some $s > 0$. Then as $\varepsilon \to 0$ the distribution of $r_{n_\varepsilon} - r_0$ converges weakly to R_s, where $R_\bullet$ is a diffusion process of the form* (2)*, with the drift and the variance*

$$b(R) = \int_0^1 E_2(\theta, R)\, d\theta, \qquad \sigma^2(R) = \int_0^1 v^2(\theta, R)\, d\theta.$$

for some function E_2, defined in (11).

- In the case that $u_{\pm 1} = v_{\pm 1}$ and they are independent of r we have two area-preserving standard maps. In this case the assumptions become
  - **[H0]** $\int v_i(\theta) d\theta = 0$ for $i = \pm 1$;
  - **[H1]** v_1 and v_{-1} have no common zeroes;
  - **[H2]** v is not identically zero.
  - **[H3]** the functions v_i are trigonometric polynomials;
  - **[H4]** the same condition as above without dependence on r;
  - **[H5]** the same condition as above without dependence on r;

A good example is $u_1(\theta) = v_1(\theta) = \cos\theta$ and $u_{-1}(\theta) = v_{-1}(\theta) = \sin\theta$. In this case

$$\int_0^1 E_2(\theta, r) d\theta \equiv 0, \qquad \sigma^2 = \int_0^1 v^2(\theta)\, d\theta$$

and for $n \le \varepsilon^{-2}$ the distribution $r_n - r_0$ converges to the zero mean variance $\varepsilon n^2\sigma^2$ normal distribution, denoted $\mathcal{N}(0, \varepsilon n^2\sigma^2)$. More generally, we have the following "vertical central limit theorem":

Theorem 2.2. *Assume that in the notations above conditions* **[H0-H5]** *hold. Let $n_\varepsilon \varepsilon^2 \to s > 0$ as $\varepsilon \to 0$ for some $s > 0$. Then as $\varepsilon \to 0$ the distribution of $r_{n_\varepsilon} - r_0$ converges weakly to a normal random variable $\mathcal{N}(0, s^2\sigma^2)$.*

- Numerical experiments of Mockel [31] show that no common fixed points [H1] (resp. [H4]) is not neccessary for Theorem 2.1 to hold. One could probably replaced by a weaker non-degeneracy condition, e.g. that the linearisation of maps $f_{\pm 1}$ at the common fixed point (resp. periodic points) are different.

- In [34] Sauzin studies random iterations of the standard maps $(\theta, r) \to (\theta + r + \lambda\phi(\theta), r + \lambda\phi(\theta))$, where λ is chosen randomly from $\{-1, 0, 1\}$ and proves the vertical central limit theorem; In [29, 35] Marco-Sauzin present examples of nearly integrable systems having a set of initial conditions exhibiting the vertical central limit theorem.

- The condition [H3] that the functions v_i are trigonometric polynomials in θ seems redundant too, however, removing it leads to considerable technical difficulties (see Section 3.2 and Remark 3.1). In short, for perturbations by a trigonometric polynomial there are finitely many resonant zones. This finiteness considerably simplifies the analysis.

- One can replace $\Sigma = \{0, 1\}^{\mathbb{Z}}$ with $\Sigma_N = \{0, 1, \ldots, N-1\}^{\mathbb{Z}}$, consider any finite number of maps of the form (4) and a transitive Markov chain with some transition probabilities. If conditions [H2–H4] are satisfied for the proper averages $\mathbb{E}v$ of v, then Theorem 2.1 holds.

3 Strategy of the proof

3.1 Strip decomposition

The main idea of the proof is to divide the cylinder $\mathbb{A}$ in strips $\mathbb{T} \times I^i_\beta$, where $I^j_\beta \subset \mathbb{R}$, $j \in \mathbb{Z}$ are intervals of size ε^β, for any $0 < \beta \le 1/5$. Then we will study how the random variable $r_n - r_0$ behaves in each strip. More precisely, decompose the process $r_n(\omega), n \in \mathbb{Z}_+$ into infinitely many time intervals defined by stopping times

$$0 < n_1 < n_2 < \ldots, \tag{7}$$

where

- $r_{n_i}(\omega)$ is ε-close to the boundary between I_β^j and I_β^{j+1} for some $j \in \mathbb{Z}$
- $r_{n_{i+1}}(\omega)$ is ε-close to the other boundary of either I_β^j or of I_β^{j+1} and $n_{i+1} > n_i$ is the smallest integer with this property.

Since $\varepsilon \ll \varepsilon^\beta$, being ε-close to the boundary of I_β^j with a negligible error means jump from I_β^j to the neighbour interval $I_\beta^{j\pm1}$. In what follows for brevity we drop dependence of $r_n(\omega)$'s on ω.

3.2 Subdivision of the cylinder into domains with different quantitative behaviour

Fix $b > 0$ such that $0 < \beta - 2b < 0.04$, small $\gamma > 0$, and $K_i := K_i(u_1, v_1, u_2, v_2)$, $i = 1, 2$, depending on functions $u_j, v_j,\ j = 1, 2$, such that $K_1 < K_2$ and all are independent of ε. Consider the ε^β-grid in $\mathbb{R}$. Denote by I_β a segment whose end points are in the grid. We distinguish among three types of strips I_β. We will have strips of three types as well as transition zones from one to another. We define:

- **The Real Rational (RR) case:** A strip I_β is called *real rational* if there exists a rational $p/q \in I_\beta$, with $\gcd(p, q) = 1$ and $|q| \leq d$. Clearly, there are just finitely many strips of this kind. However, this case is the most complicated one and requires a detailed study.
- **The Imaginary Rational (IR) case:** A strip I_β is called *imaginary rational* if there exists a rational $p/q \in I_\beta$, with $\gcd(p, q) = 1$ with $d < |q| < \varepsilon^{-b}$.

  The reason we call these strips are imaginary rational, because the leading term of the angular dynamics is a rational rotation, however, averaged systems appearing in the previous case are vanishing (see the next section).

  We show that the imaginary rational strips occupy an $\mathcal{O}(\varepsilon^\rho)$-fraction of the cylinder (see Lm. A.1 in Sect. A). We can show that orbits spend small fraction of the total time in these strips and global behaviour is determined by behaviours in the complement, which we call totally irrational.

- **The Totally Irrational (TI) case:** A strip I_β is called *totally irrational* if $r \in I_\beta$ and $|r - p/q| < \varepsilon^\beta$, with $\gcd(p,q) = 1$, then $|q| > \varepsilon^{-b}$.

  In this case, we show that there is a good "ergodization" and

  $$\sum_{k=0}^{n-1} \omega_k v\left(\theta_0 + k\frac{p}{q}\right) \approx \sum_{k=0}^{n-1} \omega_k v\left(\theta_0 + kr_0^*\right).$$

  Loosely speaking, any $r_0^* \in I_\beta \cap (\mathbb{R} \setminus \mathbb{Q})$ can be treated as an irrational. These strips cover most of the cylinder and give the dominant contribution to the behaviour of $r_n - r_0$. Eventually it will lead to the desired weak convergence to a diffusion process (Theorem 2.1).

- **Transition zones, type I:** A zone is a transition zone if there is p/q such that $\gcd(p,q) = 1$ and $|q| \le d$ and it is defined by the corresponding annuli $K_1\varepsilon^{1/2} \le |r - p/q| \le K_2\varepsilon^{1/6}$.

  Analysis in these zones needs to be adapted as "influence" of real resonances is strong.

- **Transition zones, type II:** A zone is a transition zone if there is p/q such that $\gcd(p,q) = 1$ and $|q| \le d$ and it is defined by the corresponding annuli $K_2\varepsilon^{1/6} \le |r - p/q| \le \gamma$.

  Analysis in these zones requires an adjusted coordinates, otherwise, we still study the Totally Irrational and the Imaginary Rational strips inside of the type II Transition Zones.

Remark 3.1. Notice that finiteness of Real Rational strips follows from assumption [H3]. If the expected potential is not a trigonometric polynomial in θ this is not true.

3.3 The normal form

The first step is to find a normal form, so that the deterministic part of map (6) is as simple as possible. In short, we shall see that the deterministic system in both the Totally Irrational case and the Imaginary rational case are a small perturbation of the perfect twist map:

$$\begin{pmatrix} \theta \\ r \end{pmatrix} \longmapsto \begin{pmatrix} \theta + r \\ r \end{pmatrix}.$$

On the contrary, in the Real Rational case, the deterministic system will be close to a pendulum-like system:

$$\begin{pmatrix} \theta \\ r \end{pmatrix} \longmapsto \begin{pmatrix} \theta + r \\ r + \varepsilon E(\theta, r) \end{pmatrix},$$

for an "averaged" potential $E(\theta, r)$ (see e.g. Thm. 4.2, (12)). We note that this system has the following approximate first integral:

$$H(\theta, r) = \frac{r^2}{2} - \varepsilon \int_0^\theta E(s, r) ds, \tag{8}$$

so that indeed it is close to a pendulum-like system. This will lead to different qualitative behaviours when considering the random system. Inside the Real Rational strips as well as the transition zones we use H as one of the coordinates.

The rigorous statement of these results about the normal forms is given in Theorem 4.2, Sect. 4.

3.4 Analsys of the Martingale problem in each kind of strip

The next step is to study the behaviour of the random system respectively in Totally Irrational, Imaginary Rational and Real Rational strips, as well as in the Transition Zones. This is done in Sections 5.1–5.4. More precisely, we use a discrete version of the scheme by Freidlin and Wentzell [20], giving a sufficient condition to have weak convergence to a diffusion process as $\varepsilon \to 0$ in terms of the associated Martingale problem (see Lemma B.1). Now using the results proved below we derive the main result — Theorem 2.1. This is done in two steps. First, we describe local behaviour in each strip and then we combine the information. Fix $s > 0$.

By the discrete version of Lemma B.1 is sufficient to prove that as $\varepsilon \to 0$ any time $n \leq s\varepsilon^{-2}$ and any (θ_0, r_0) we have

$$\begin{aligned} \mathbb{E} \Big(& e^{-\lambda\varepsilon^2 n} f(r_n) + \\ & \varepsilon^2 \sum_{k=0}^{n-1} e^{-\lambda\varepsilon^2 k} \quad \Big[\lambda f(r_k) - \Big(b(r_k) f'(r_k) + \frac{\sigma^2(r_k)}{2} f''(r_k) \Big) \Big] \Big) - f(r_0) \to 0, \end{aligned} \tag{9}$$

We define Markov times $0 = n_0 < n_1 < n_2 < \cdots < n_{m-1} < n_m < n$ for some random $m = m(\omega)$ such that each n_k is the stopping time as in (7). Almost surely $m(\omega)$ is finite. We decompose the above sum

$$\begin{aligned} \sum_{k=0}^{m} \quad & \mathbb{E} \Big(e^{-\lambda\varepsilon^2 n_{k+1}} f(r_{n_{k+1}}) - e^{-\lambda\varepsilon^2 n_k} f(r_{n_k}) + \\ & \varepsilon^2 \sum_{s=n_k}^{n_{k+1}} e^{-\lambda\varepsilon^2 s} \Big[\lambda f(r_s) - \Big(b(r_s) f'(r_s) + \frac{\sigma^2(r_s)}{2} f''(r_s) \Big) \Big] \Big) \end{aligned}$$

and show that it converges to $f(r_0)$.

3.4.1 A Totally Irrational Strip

Let the drift and the variance be

$$b(r) = \int_0^1 E_2(\theta, r)\, d\theta \quad \text{and} \quad \sigma^2(r) = \int_0^1 v^2(\theta, r)\, d\theta,$$

where the function E_2 is defined in (11). Let r_0 be ε-close to the boundary of two totally irrational strips and let n_β be stopping of hitting ε-neighbourhoods of the adjacent boundaries. In Lemma 5.3 we prove that

$$\begin{aligned}\mathbb{E}\Bigg(&e^{-\lambda\varepsilon^2 n_\beta} f(r_{n_\beta}) + \\ &\varepsilon^2 \sum_{k=0}^{n_\beta - 1} e^{-\lambda\varepsilon^2 k}\left[\lambda f(r_k) - \left(b(r_k) f'(r_k) + \frac{\sigma^2(r_k)}{2} f''(r_k)\right)\right]\Bigg) \\ &\qquad\qquad -f(r_0) = \mathcal{O}(\varepsilon^{2\beta+d}),\end{aligned}$$

for some $d > 0$.

3.4.2 An Imaginary Rational Strip

Let the drift and the variance be

$$b_{IR}(\theta, r) = \frac{1}{q}\sum_{k=0}^{q-1} E_2(\theta + kr, r) \quad \text{and} \quad \sigma^2_{IR}(\theta, r) = \frac{1}{q}\sum_{k=0}^{q-1} v^2(\theta + kr, r).$$

Let r_0 be ε-close to the boundary of an imaginary rational strip and let n_β be stopping of hitting ε-neighbourhoods of the adjacent boundaries. In Lemma 5.5 we prove that

$$\begin{aligned}\mathbb{E}\Bigg(&e^{-\lambda\varepsilon^2 n_\beta} f(r_{n_\beta}) + \\ &\varepsilon^2 \sum_{k=0}^{n_\beta - 1} e^{-\lambda\varepsilon^2 k}\left[\lambda f(r_k) - \left(b(\theta_k, r_k) f'(r_k) + \frac{\sigma^2(\theta_k, r_k)}{2} f''(r_k)\right)\right]\Bigg) \\ &\qquad\qquad -f(r_0) = \mathcal{O}(\varepsilon^{2\beta+d}),\end{aligned}$$

As one can see, the limiting process does not take place on a line, since the drift and diffusion coefficient depend also on the variable θ.

Notice that the drift $b(\theta, r)$ and the variance $\sigma(\theta, r)$ both are θ-dependent functions. In Section 3.5 we show that time spent in these strips is too small to affect the drift and the variance of the limiting process.

3.4.3 A Real Rational Strip

Let in the rescaled variable $r - p/q = R\sqrt{\varepsilon}$ the drift and the variance be

$$b_{RR}(\theta, R) = F(\theta, R), \qquad \sigma^2_{RR}(\theta, R) = (Rp/q)^2 \sum_{k=0}^{q-1} v^2(\theta + kR, R),$$

where F is some function to be defined in (92). Consider the Real Rational case assuming that

$$|r - p/q| \le K_1 \varepsilon^{1/2} \qquad \Longleftrightarrow \qquad |R| \le K_1$$

that is, that r is close to the "pendulum" domain. In this case, we study the process (θ_{qn}, H_n) with $H_n := H^{p/q}(\theta_{qn}, R_{qn})$, where $H^{p/q}(\theta, R)$ is an approximate first integral of the deterministic system (8). In the rescaled variables it has the form

$$H^{p/q}(\theta, R) = \frac{R^2}{2} - V^{p/q}(\theta),$$

where

$$V^{p/q}(\theta) = \int_0^\theta \mathbb{E} v_{p,q}(s, p/q)\, ds$$

for a properly defined averaged potential (see Thm. 4.2, (14)). In Lemma 5.6 we prove that, $H_n - H_0$ converges weakly to a diffusion process R_t with $t = \varepsilon^2 n$.

Notice that the limiting process does not take place on a line. In this case it takes place on a graph, similarly as in [20]. More precisely, consider the level sets of the function $H^{p/q}(\theta, R)$. The critical points of the potential $V^{p/q}(\theta)$ give rise to critical points of the associated Hamiltonian system. Moreover, if the critical point is a local minimum of V, then it corresponds to a focus of the Hamiltonian system, while if it is a local maximum of $V^{p/q}$, then it corresponds to a saddle of the Hamiltonian system. Now, if for every value $H \in \mathbb{R}$ we identify all the points (θ, R) in the same connected component of the curve $\{H^{p/q}(\theta, R) = H\}$, we obtain a graph Γ (see Figure 2 for an example). The interior vertices of this graph represent the saddle points of the underlying Hamiltonian system jointly with their separatrices, while the exterior vertices represent the focuses of the underlying Hamiltonian system. Finally, the edges of the graph represent the domains that have the separatrices as boundaries. The process H_n takes places on this graph, and so it is a diffusion process on a graph.

3.4.4 A transition zone

Finally, in Lemma 5.7 we deal with the Transition Zones of Type I and Type II, that is the zones in the Real Rational strips such that $K_1 \le |R| \le K_2 \varepsilon^{-1/3}$



(a) (b)

Figure 2: (a) A potential and the phase portrait of its corresponding Hamiltonian system. (b) The associated graph Γ.

and $K_2\varepsilon^{-1/3} \le |R| \le \gamma\varepsilon^{-1/2}$. In these strips we study the process $(\theta_{nq}, H_{nq}) = (\theta_{nq}, H(\theta_{qn}, R_{qn}))$. In this regime we fix small $\rho > 0$ and subdivide each zone in sub-strips

$$I_\rho(R_0) = \{H \in \mathbb{R} \,:\, |H - H_0| \le |R_0|\, \varepsilon^{1/2-\rho}\}.$$

We prove that, inside each of one these sub-strips, as $\varepsilon \to 0$ the process $H_n - H_0$ converges weakly to a diffusion process R_t with $t = \varepsilon^2 n$, zero drift and the variance:

$$\sigma^2_{TZ}(\theta, R) = |R|^2 \sum_{k=0}^{q-1} v^2(\theta + kR, R).$$

3.5 From the local diffusion in the rational strips to the global diffusion on the line

In this section we resolve the following problem. In order to combine all the previous results, which characterise the local behaviour of the process inside of infinitesimally small strips, to determine the global behaviour of the process in a $\mathcal{O}(1)-$strip. First, we prove that the Imaginary Rational and Real Rational strips cover a negligibly small part of any $\mathcal{O}(1)-$strip (see Section A). Then, one can argue that the process is determined by the process in the Totally Irrational strips.

Notice both the drift $b_{IR}(\theta, r)$ and the variance $b_{IR}(\theta, r)$ at any Imaginary Rational strip is given by θ-dependent functions. Our main result (Theorem 2.1), however, is a diffusion process on a line. To prove that this dependence does not enter into the global diffusion process we show that the process spends infinitesimal amount of time inside of those strips as follows.

Lemma 3.2. *Let* $(\theta_k, r_k) = f^k_{\Omega_k}(\theta_0, r_0), k \geq 1$ *be a random orbit defined by (5) for some random sequence* $\{\omega_k\}_{k\in\mathbb{Z}_+}$. *Let* $n \leq \varepsilon^{-2}$ *and*

$$T_R(n) = \#\{0 \leq k \leq n : r_k \text{ belongs to either}$$

$$\text{an Imaginary Rational or a Real Rational strip}\}.$$

Then for any $\rho > 0$ *and* $\varepsilon > 0$ *small enough*

$$\mathbb{P}\{T_R(n) \geq \rho n\} \leq \rho.$$

Proof. Define

$$b_{IR}(r) := \min_{\theta\in[0,1)} b_{IR}(\theta, r) \quad \text{and} \quad \sigma^2_{IR}(r) := \min_{\theta\in[0,1)} \sigma^2_{IR}(\theta, r)$$

Consider the process R^{IR}_t with the drift $b_{IR}(r)$ and the variance $\sigma^2_{IR}(r)$. By definition this process spends more time in I_β that the process with the drift $b_{IR}(\theta, r)$ and the variance $\sigma^2_{IR}(\theta, r)$. Moreover, it is a diffusion process on a line. Then, using a local time argument, it can be seen that the time spent on a given domain is proportional to the size of this domain up to a uniform constant. Hence, the time the original process spends in all the Imaginary Rational strips is infinitesimally small compared to the time it spends on the Totally Irrational ones. However, the time spent in the Imaginary Rational strip could be infinite and the argument would not be valid. This cannot happen, since if r belongs to an Imaginary Rational strip one has that $\sigma^2(\theta, r) \neq 0$. Thus, it is enough to prove that for all imaginary rational p/q one has $\sigma^2(\theta, p/q) \neq 0$. Indeed, if this is true, then for $|r - p/q| \leq \varepsilon^\beta$ and ε is sufficiently small, one has that $\sigma^2(\theta, r) \neq 0$ by Lemma 3.3.

Finally, in the Real Rational case one can use a result from [19] that diffusion processes on a graph have well-defined local time. Thus, the time spent in all the Real Rational strips is infinitesimally small compared to the time spent in the Totally Irrational ones. Now one can have $\sigma^2_{RR} = 0$, but it happens just when $r = p/q$, which follows directly from assumption [**H2**]. In this case, one can see that $b_{RR}(\theta, r) \neq 0$, so that the process is non-degenerate and thus the fraction of time spend in the Real Rational strips is less than any ahead given fraction. □

Lemma 3.3. $\sigma^2(\theta, p/q) \neq 0$ *if* p/q *is any Imaginary Rational.*

Proof. On the one hand, if $d < |q| \leq 2d$ this is ensured by hypothesis [**H4**]. On the other hand, if $|q| > 2d$ then $\sigma^2(\theta, p/q) = 0$ implies:

$$v(\theta + kp/q, p/q) = 0, \qquad k = 0, \cdots, q-1. \tag{10}$$

Now, since $v(\theta, p/q)$ is a trigonometric polynomial in θ of degree d, it can have at most $2d$ zeros, or else be identically equal to zero. The latter case cannot occur, since by assumption **[H2]** we know that

$$\int_0^1 v^2(\theta, r) \neq 0 \quad \text{for all} \quad r \in \mathbb{R},$$

so that $v(\theta, p/q) \not\equiv 0$. Thus, $v(\theta, p/q)$ has at most $2d$ zeros. Consequently equation (10) cannot be satisfied for all $k = 0, \cdots, q-1$, since $|q| > 2d$, so that $\sigma^2(\theta, p/q) \neq 0$. The same argument applies to the Transition Zones. $\square$

Combining these facts one can apply the arguments from [20], sect. 8 and prove that the limiting diffusion process has the drift $b(r)$ and the variance $\sigma^2(r)$ corresponding to Totally Irrational strips.

3.6 Plan of the rest of the paper

In Section 4 we state and prove the normal form theorem for the expected cylinder map $\mathbb{E}f$. Main difference with a typical normal form is that we need to have not only the leading term in ε, but also ε^2-terms. The latter terms give information about the drift $b(r)$ (see (11)).

In Section 5.1 we analyse the Totally Irrational case and prove approximation for the expectation from Section 3.4.1.

In Section 5.2 we analyse the Imaginary Rational case and prove an analogous formula from Section 3.4.2.

In Section 5.3 we analyse the Real Rational case and prove an analogous formula from Section 3.4.3.

In Section 5.4 we study the Transition Zones and prove an analogous formula from Section 3.4.4.

In Section A we estimate measure of the complement to the Totally Irrational strips and the Transition Zones of type II.

In Section B we present several auxiliary lemmas used in the proof.

4 The Normal Form Theorem

In this section we shall prove the Normal Form Theorem, which will allow us to deal with the simplest possible deterministic system. To this end, we shall enunciate a technical lemma which we will need in the proof of the theorem. This is a simplified version (sufficient for our purposes) of Lemma 3.1 in [5].

Lemma 4.1. *Let $g(\theta, r) \in \mathcal{C}^l(\mathbb{T} \times B)$, where $B \subset \mathbb{R}$. Then:*

1. *If $l_0 \leq l$ and $k \neq 0$, $\|g_k(r) e^{2\pi i k\theta}\|_{\mathcal{C}^{l_0}} \leq |k|^{l_0 - l} \|g\|_{\mathcal{C}^{l_0}}$.*

2. *Let $g_k(r)$ be some functions that satisfy $\|\partial_{r^\alpha} g_k\|_{\mathcal{C}^0} \leq M|k|^{-\alpha-2}$ for all $\alpha \leq l_0$ and some $M > 0$. Then:*
$$\left\| \sum_{\substack{k\in\mathbb{Z} \\ 0<k\leq d}} g_k(r) e^{2\pi i k\theta} \right\|_{\mathcal{C}^{l_0}} \leq cM,$$
for some constant c depending on l_0.

Let $\mathcal{R}$ be the finite set of resonances of this map, namely,

$$\mathcal{R} = \{p/q \in \mathbb{Q} \,:\, \gcd(p,q) = 1, |q| < d\}.$$

Denote by $\mathcal{O}(\varepsilon)$ a function whose $\mathcal{C}^0$-norm is bounded by $C\varepsilon$ for some C independent of ε.

Define

$$E_2(\theta, r) = \mathbb{E}v(\theta, r)\, \partial_\theta S_1(\theta, r) + \mathbb{E}w(\theta, r), \quad b(r) = \int E_2(\theta, r) d\theta, \tag{11}$$

where S_1 is a certain generating function defined in (21–22).

Theorem 4.2. *Consider the expected map $\mathbb{E}f$ of the map* (4)

$$\mathbb{E}f \begin{pmatrix} \theta \\ r \end{pmatrix} \longmapsto \begin{pmatrix} \theta + r + \varepsilon \mathbb{E}u(\theta, r) + \mathcal{O}_l(\varepsilon^{1+a}) \\ r + \varepsilon \mathbb{E}v(\theta, r) + \varepsilon^2 \mathbb{E}w(\theta, r) + \mathcal{O}_l(\varepsilon^{2+a}) \end{pmatrix}.$$

Assume that the functions $\mathbb{E}u(\theta, r)$, $\mathbb{E}v(\theta, r)$ and $\mathbb{E}w(\theta, r)$ are $\mathcal{C}^l$, $l \geq 3$ and $\gamma > 0$ small. Then there exists $K > 0$ independent of ε and a canonical change of variables:

$$\begin{array}{rcl} \Phi : \mathbb{T} \times \mathbb{R} & \to & \mathbb{T} \times \mathbb{R}, \\ (\tilde\theta, \tilde r) & \mapsto & (\theta, r), \end{array}$$

such that:

- *If $|\tilde r - p/q| \geq \gamma$ for all $p/q \in \mathcal{R}$, then:*
$$\begin{aligned} &\Phi^{-1} \circ \mathbb{E}f \circ \Phi(\tilde\theta, \tilde r) = \\ &\qquad \begin{pmatrix} \tilde\theta + \tilde r + \varepsilon \mathbb{E}u(\theta, r) + \varepsilon E_1(\theta, r) + \mathcal{O}_l(\varepsilon^{1+a}) \\ \tilde r + \varepsilon^2 E_2(\tilde\theta, \tilde r) + \mathcal{O}_l(\varepsilon^{2+a}) \end{pmatrix}, \end{aligned} \tag{12}$$
  where E_1 and E_2 are some $\mathcal{C}^{l-1}$ functions. Moreover, E_2 verifies $\|E_2\|_{\mathcal{C}^0} \leq K$ and it verifies:
$$\begin{aligned} &b(r) := \int_0^1 E_2(\tilde\theta, \tilde r) d\tilde\theta = \\ &\int_0^1 \left[\partial_r \mathbb{E}v(\tilde\theta, \tilde r) \partial_\theta S_1(\tilde\theta, \tilde r) - \partial_\theta^2 S_1(\tilde\theta, \tilde r) \left(\mathbb{E}u(\tilde\theta, \tilde r) - \mathbb{E}v(\tilde\theta, \tilde r) \right) \right] d\tilde\theta. \end{aligned} \tag{13}$$
  In particular, $b(r)$ satisfies $\|b\|_{\mathcal{C}^0} \leq K$ and in the area-preserving case (when $\mathbb{E}u(\theta, r) = \mathbb{E}v(\theta, r) = \mathbb{E}v(\theta)$), $b(r) \equiv 0$.

- *If* $|\tilde{r} - p/q| \leq 2\gamma$ *for a given* $p/q \in \mathcal{R}$*, then:*

$$\Phi^{-1} \circ \mathbb{E}f \circ \Phi(\tilde{\theta}, \tilde{r}) = \tag{14}$$

$$\begin{pmatrix} \tilde{\theta} + \tilde{r} + \varepsilon \left[\mathbb{E}u(\tilde{\theta}, p/q) - \mathbb{E}v(\tilde{\theta}, p/q) + \mathbb{E}v_{p,q}(\tilde{\theta}, p/q) + E_3(\tilde{\theta})\right] + \mathcal{O}_l(\varepsilon^{1+a}) \\ \tilde{r} + \varepsilon \mathbb{E}v_{p,q}(\tilde{\theta}, \tilde{r}) + \varepsilon^2 E_4(\tilde{\theta}, \tilde{r}) + \mathcal{O}_l(\varepsilon^{2+a}) \end{pmatrix},$$

where $\mathbb{E}v_{p,q}$ *is the* $\mathcal{C}^l$ *function defined as:*

$$\mathbb{E}v_{p,q}(\tilde{\theta}, \tilde{r}) = \sum_{k \in \mathcal{R}_{p,q}} \mathbb{E}v^k(\tilde{r}) e^{2\pi i k \tilde{\theta}},$$

and E_3 *is the* $\mathcal{C}^{l-1}$ *function:*

$$E_3(\tilde{\theta}) = -\sum_{k \notin \mathcal{R}_{p,q}} \frac{i(\mathbb{E}v^k)'(p/q)}{2\pi k} e^{2\pi i k \tilde{\theta}},$$

where $\mathcal{R}_{p,q} = \{k \in \mathbb{Z} \,:\, k \neq 0,\ |k| < d,\ kp/q \in \mathbb{Z}\}$*.*

Moreover, E_4 *is a* $\mathcal{C}^{l-1}$ *function verifying* $\|E_4\|_{\mathcal{C}^0} \leq K$*.*

Also, Φ *is* $\mathcal{C}^2$*-close to the identity. More precisely, there exists a constant* M *independent of* ε *such that:*

$$\|\Phi - \mathrm{Id}\|_{\mathcal{C}^2} \leq M\varepsilon. \tag{15}$$

Proof. For each $p/q \in \mathcal{R}$ we will perform a different change. Since the procedure is the same for all $p/q \in \mathcal{R}$, from now on we fix $p/q \in \mathcal{R}$. The procedure for the rest is analogous.

We will consider the canonical change defined implicitly by a given generating function $S(\theta, \tilde{r}) = \theta\tilde{r} + \varepsilon S_1(\theta, \tilde{r})$, that is:

$$\begin{aligned} r &= \partial_\theta S(\theta, \tilde{r}) = \tilde{r} + \varepsilon \partial_\theta S_1(\theta, \tilde{r}) \\ \tilde{\theta} &= \partial_{\tilde{r}} S(\theta, \tilde{r}) = \theta + \varepsilon \partial_{\tilde{r}} S_1(\theta, \tilde{r}). \end{aligned}$$

We shall start by writing explicitly the first orders of the ε-series of $\Phi^{-1} \circ \mathbb{E}f \circ \Phi$. If $(\theta, r) = \Phi(\tilde{\theta}, \tilde{r})$ is the change given by the generating function S, then one has:

$$\Phi(\tilde{\theta}, \tilde{r}) =$$

$$\begin{pmatrix} \tilde{\theta} - \varepsilon \partial_{\tilde{r}} S_1(\tilde{\theta}, \tilde{r}) + \varepsilon^2 \partial_\theta \partial_{\tilde{r}} S_1(\tilde{\theta}, \tilde{r}) \partial_{\tilde{r}} S_1(\tilde{\theta}, \tilde{r}) + \mathcal{O}(\varepsilon^3 \|\partial_\theta^2 \partial_{\tilde{r}} S_1 (\partial_{\tilde{r}} S_1)^2\|_{\mathcal{C}^0}) \\ \tilde{r} + \varepsilon \partial_\theta S_1(\tilde{\theta}, \tilde{r}) - \varepsilon^2 \partial_\theta^2 S_1(\tilde{\theta}, \tilde{r}) \partial_{\tilde{r}} S_1(\tilde{\theta}, \tilde{r}) + \mathcal{O}(\varepsilon^3 \|\partial_\theta^3 S_1 (\partial_{\tilde{r}} S_1)^2\|_{\mathcal{C}^0}) \end{pmatrix}. \tag{16}$$

Its inverse is given by:

$$\Phi^{-1}(\theta, r) =$$

$$\begin{pmatrix} \theta + \varepsilon \partial_{\tilde{r}} S_1(\theta, r) - \varepsilon^2 \partial_{\tilde{r}}^2 S_1(\theta, r) \partial_\theta S_1(\theta, r) + \mathcal{O}(\varepsilon^3 \|\partial_{\tilde{r}}^3 S_1 (\partial_\theta S_1)^2\|_{\mathcal{C}^0}) \\ r - \varepsilon \partial_\theta S_1(\theta, r) + \varepsilon^2 \partial_\theta \partial_{\tilde{r}} S_1(\theta, r) \partial_\theta S_1(\theta, r) + \mathcal{O}(\varepsilon^3 \|\partial_\theta \partial_{\tilde{r}}^2 S_1 (\partial_\theta S_1)^2\|_{\mathcal{C}^0}) \end{pmatrix}. \tag{17}$$

Now, first we compute $\mathbb{E}f \circ \Phi(\tilde{\theta}, \tilde{r})$. One can see that:

$$\mathbb{E}f \circ \Phi(\tilde{\theta}, \tilde{r}) = \begin{pmatrix} \tilde{\theta} + r\varepsilon A_1 + \varepsilon^2 A_2 + \varepsilon^3 A_3 \\ r + \varepsilon B_1 + \varepsilon^2 B_2 + \varepsilon^3 B_3 \end{pmatrix},$$

where:

$$\begin{aligned}
A_1 &= \mathbb{E}u(\tilde{\theta}, \tilde{r}) - \partial_{\tilde{r}} S_1(\tilde{\theta}, \tilde{r}) + \partial_\theta S_1(\tilde{\theta}, \tilde{r}) \\
A_2 &= -\partial_\theta \mathbb{E}u(\tilde{\theta}, \tilde{r}) \partial_{\tilde{r}} S_1(\tilde{\theta}, \tilde{r}) + \partial_r \mathbb{E}u(\tilde{\theta}, \tilde{r}) \partial_\theta S_1(\tilde{\theta}, \tilde{r}) + \partial_\theta \partial_{\tilde{r}} S_1(\tilde{\theta}, \tilde{r}) \partial_{\tilde{r}} S_1(\tilde{\theta}, \tilde{r}) \\
&\quad -\partial_\theta^2 S_1(\tilde{\theta}, \tilde{r}) \partial_{\tilde{r}} S_1(\tilde{\theta}, \tilde{r}), \\
A_3 &= \mathcal{O}(\|\partial_\theta^2 \partial_{\tilde{r}} S_1 (\partial_{\tilde{r}} S_1)^2\|_{\mathcal{C}^0}) + \mathcal{O}(\|\partial_\theta^3 S_1 (\partial_{\tilde{r}} S_1)^2\|_{\mathcal{C}^0}) \\
&\quad +\mathcal{O}(\|\mathbb{E}u\|_{\mathcal{C}^1} \|\partial_\theta S_1\|_{\mathcal{C}^1} \|\partial_{\tilde{r}} S_1\|_{\mathcal{C}^0}) + \mathcal{O}(\|\mathbb{E}u\|_{\mathcal{C}^2} (\|\partial_\theta S_1\|_{\mathcal{C}^0} + \|\partial_{\tilde{r}} S_1\|_{\mathcal{C}^0})^2),
\end{aligned}$$

and:

$$\begin{aligned}
B_1 &= \mathbb{E}v(\tilde{\theta}, \tilde{r}) + \partial_\theta S_1(\tilde{\theta}, \tilde{r}), \\
B_2 &= -\partial_\theta \mathbb{E}v(\tilde{\theta}, \tilde{r}) \partial_{\tilde{r}} S_1(\tilde{\theta}, \tilde{r}) + \partial_r \mathbb{E}v(\tilde{\theta}, \tilde{r}) \partial_\theta S_1(\tilde{\theta}, \tilde{r}) \\
&\quad -\partial_\theta^2 S_1(\tilde{\theta}, \tilde{r}) \partial_{\tilde{r}} S_1(\tilde{\theta}, \tilde{r}), \\
B_3 &= \mathcal{O}(\|\partial_\theta^3 S_1 (\partial_{\tilde{r}} S_1)^2\|_{\mathcal{C}^0}) + \mathcal{O}(\|\mathbb{E}v\|_{\mathcal{C}^1} \|\partial_\theta S_1\|_{\mathcal{C}^1} \|\partial_{\tilde{r}} S_1\|_{\mathcal{C}^0}) \\
&\quad +\mathcal{O}(\|\mathbb{E}v\|_{\mathcal{C}^2} (\|\partial_\theta S_1\|_{\mathcal{C}^0} + \|\partial_{\tilde{r}} S_1\|_{\mathcal{C}^0})^2).
\end{aligned} \tag{18}$$

Then, using (17) one can see that:

$$\Phi^{-1} \circ \mathbb{E}f \circ \Phi(\tilde{\theta}, \tilde{r}) = \begin{pmatrix} \tilde{\theta} + r + \varepsilon \hat{A}_1 + \varepsilon^2 \hat{A}_2 \\ r + \varepsilon \hat{B}_1 + \varepsilon^2 \hat{B}_2 + \varepsilon^3 \hat{B}_3 \end{pmatrix}, \tag{19}$$

where:

$$\begin{aligned}
\hat{A}_1 &= A_1 + \partial_{\tilde{r}} S_1(\tilde{\theta} + \tilde{r}, \tilde{r}), \\
\hat{A}_2 &= A_2 + \varepsilon A_3 + \mathcal{O}(\|\partial_\theta \partial_{\tilde{r}} S_1 A_1\|_{\mathcal{C}^0}) + \mathcal{O}(\|\partial_{\tilde{r}}^2 S_1 B_1\|_{\mathcal{C}^0}) \\
&\quad +\mathcal{O}(\|\partial_{\tilde{r}}^2 S_1 \partial_\theta S_1\|_{\mathcal{C}^0}),
\end{aligned}$$

and:

$$\begin{aligned}
\hat{B}_1 &= B_1 - \partial_\theta S_1(\tilde{\theta} + \tilde{r}, \tilde{r}) \\
\hat{B}_2 &= B_2 - \partial_\theta^2 S_1(\tilde{\theta} + \tilde{r}, \tilde{r}) A_1 - \partial_{\tilde{r}} \partial_\theta S_1(\tilde{\theta} + \tilde{r}, \tilde{r}) B_1 \\
&\quad +\partial_\theta \partial_{\tilde{r}} S_1(\tilde{\theta} + \tilde{r}, \tilde{r}) \partial_\theta S_1(\tilde{\theta} + \tilde{r}, \tilde{r}), \\
\hat{B}_3 &= B_3 + \mathcal{O}(\|\partial_\theta \partial_{\tilde{r}}^2 S_1 (\partial_\theta S_1)^2\|_{\mathcal{C}^0}) \\
&\quad +\mathcal{O}(\|\partial_\theta^2 S_1 (A_2 + \varepsilon A_3)\|_{\mathcal{C}^0} + \|\partial_\theta \partial_{\tilde{r}} S_1 B_2\|_{\mathcal{C}^0}) \\
&\quad +\mathcal{O}(\|\partial_\theta^3 S_1 A_1^2\|_{\mathcal{C}^0} + \|\partial_\theta^2 \partial_{\tilde{r}} S_1 A_1 B_1\|_{\mathcal{C}^0} + \|\partial_\theta \partial_{\tilde{r}}^2 S_1 B_1^2\|_{\mathcal{C}^0}) \\
&\quad +\mathcal{O}(\|\partial_\theta^2 \partial_{\tilde{r}} S_1 A_1 \partial_\theta S_1\|_{\mathcal{C}^0} + \|\partial_\theta \partial_{\tilde{r}}^2 S_1 B_1 \partial_\theta S_1\|_{\mathcal{C}^0}) \\
&\quad +\mathcal{O}(\|\partial_\theta \partial_{\tilde{r}} S_1 \partial_\theta^2 S_1 A_1\|_{\mathcal{C}^0} + \|(\partial_\theta \partial_{\tilde{r}} S_1)^2 B_1\|_{\mathcal{C}^0}).
\end{aligned} \tag{20}$$

Now that we know the terms of order ε and ε^2 of $\Phi^{-1} \circ \mathbb{E}f \circ \Phi$, we shall proceed to find a suitable $S_1(\theta, \tilde{r})$ such that these terms are as simple as possible. More precisely, we want to simplify the second component of (19). Ideally we would like that $\hat{B}_1 = 0$. Namely, we want to solve the following equation whenever it is possible:

$$\partial_\theta S_1(\tilde{\theta}, \tilde{r}) + \mathbb{E}v(\tilde{\theta}, \tilde{r}) - \partial_\theta S_1(\tilde{\theta} + \tilde{r}, \tilde{r}) = 0.$$

One can easily find a solution of this equation by solving the corresponding equation for the Fourier coefficients. To that aim, we write S_1 and $\mathbb{E}v$ in their Fourier series:

$$S_1(\theta, \tilde{r}) = \sum_{k \in \mathbb{Z}} S_1^k(\tilde{r}) e^{2\pi i k\theta}, \tag{21}$$

$$\mathbb{E}v(\theta, r) = \sum_{\substack{k \in \mathbb{Z} \\ 0<|k|\leq d}} \mathbb{E}v^k(r) e^{2\pi i k\theta}.$$

It is obvious that for $k > d$ and $k = 0$ we can take $S_1^k(\tilde{r}) = 0$. For $0 < k \leq d$ we obtain the following homological equation for $S_1^k(\tilde{r})$:

$$2\pi i k S_1^k(\tilde{r}) \left(1 - e^{2\pi i k\tilde{r}}\right) + \mathbb{E}v^k(r) = 0. \tag{22}$$

Clearly, this equation cannot be solved if $e^{2\pi i k\tilde{r}} = 1$, i.e. if $k\tilde{r} \in \mathbb{Z}$. We note that there exists a constant L, independent of ε, $L < d^{-1}$, such that for all $0 < k \leq d$, if $\tilde{r} \neq p/q$ satisfies:

$$0 < |\tilde{r} - p/q| \leq L$$

then $k\tilde{r} \notin \mathbb{Z}$. Thus, restricting ourselves to the domain $|\tilde{r} - p/q| \leq L$, we have that if $kp/q \notin \mathbb{Z}$ equation (22) always has a solution, and if $kp/q \in \mathbb{Z}$ this equation has a solution except at $\tilde{r} = p/q$. Moreover, in the case that the solution exists, it is equal to:

$$S_1^k(\tilde{r}) = \frac{i\mathbb{E}v^k(r)}{2\pi k \left(1 - e^{2\pi i k\tilde{r}}\right)}.$$

We will modify this solution slightly to make it well defined also at $\tilde{r} = p/q$. To this end, let us consider a $\mathcal{C}^\infty$ function $\mu(x)$ such that:

$$\mu(x) = \begin{cases} 1 & \text{if} \quad |x| \leq 1, \\ 0 & \text{if} \quad |x| \geq 2, \end{cases}$$

and $0 < \mu(x) < 1$ if $x \in (1, 2)$. Then we define:

$$\mu_k(\tilde{r}) = \mu\left(\frac{1 - e^{2\pi i k\tilde{r}}}{2\pi k\gamma}\right),$$

and take:

$$S_1^k(\tilde{r}) = \frac{i\mathbb{E}v^k(r)(1-\mu_k(\tilde{r}))}{2\pi k(1-e^{2\pi i k\tilde{r}})}. \tag{23}$$

We note that this function is well defined since the numerator is identically zero in a neighbourhood of $\tilde{r} = p/q$, the unique zero of the denominator (if it is a zero indeed, that is, if $k \in \mathcal{R}_{p,q}$). More precisely, we claim that:

$$\mu_k(\tilde{r}) = \begin{cases} 1 & \text{if} \quad k \in \mathcal{R}_{p,q} \text{ and } |\tilde{r}-p/q| \leq \gamma/2, \\ 0 & \text{if} \quad k \in \mathcal{R}_{p,q} \text{ and } |\tilde{r}-p/q| \geq 3\gamma, \\ 0 & \text{if} \quad k \notin \mathcal{R}_{p,q}. \end{cases} \tag{24}$$

Indeed if $k \in \mathcal{R}_{p,q}$ there exists a constant M independent of $\tilde{r}$ and ε such that:

$$\frac{1}{\gamma}|\tilde{r}-p/q|(1-M|\tilde{r}-p/q|) \leq \left|\frac{1-e^{2\pi i k\tilde{r}}}{2\pi k\gamma}\right| \leq \frac{1}{\gamma}|\tilde{r}-p/q|(1+M|\tilde{r}-p/q|).$$

Then, on the one hand, if $k \in \mathcal{R}_{p,q}$ and $|\tilde{r}-p/q| \leq \gamma/2$ we have:

$$\left|\frac{1-e^{2\pi i k\tilde{r}}}{2\pi k\gamma}\right| \leq \frac{1}{2} + \frac{M}{4}\gamma < 1,$$

for ε sufficiently small, and thus $\mu_k(\tilde{r}) = 1$. On the other hand, if $|\tilde{r}-p/q| \geq 3\gamma$ then:

$$\left|\frac{1-e^{2\pi i k\tilde{r}}}{2\pi k\gamma}\right| \geq 3 - 9M\gamma > 2,$$

for ε sufficiently small, and thus $\mu_k(\tilde{r}) = 0$. Finally, if $k \notin \mathcal{R}_{p,q}$ then:

$$\left|\frac{1-e^{2\pi i k\tilde{r}}}{2\pi k\gamma}\right| \geq M\gamma > 2$$

for γ sufficiently small and then we also have $\mu_k(\tilde{r}) = 0$.

Now we proceed to check that the first order terms of (19) take the form (12) if $|\tilde{r}-p/q| \geq 3\gamma$ and (14) if $|\tilde{r}-p/q| \leq \gamma$. On the one hand, by definitions (23) of the coefficients $S_1^k(\tilde{r})$ and (20) of $\hat{B}_1$, we have:

$$\hat{B}_1 = \sum_{0<|k|\leq d} \mu_k(\tilde{r})\mathbb{E}v^k(\tilde{r})e^{2\pi i k\tilde{\theta}}.$$

Then, recalling (24) we obtain:

$$\hat{B}_1 = \begin{cases} 0 & \text{if} \quad |\tilde{r}-p/q| \geq \gamma \\ \displaystyle\sum_{k\in\mathcal{R}_{p,q}} \mathbb{E}v^k(\tilde{r})e^{2\pi i k\tilde{\theta}} = \mathbb{E}v_{p,q}(\tilde{\theta},\tilde{r}) & \text{if} \quad |\tilde{r}-p/q| \leq \gamma/2. \end{cases} \tag{25}$$

where we have used the definition (15) of $\mathbb{E}v_{p,q}(\tilde{\theta},\tilde{r})$. On the other hand, from the definition (23) of $S_1^k(\tilde{r})$ one can check that:

$$\begin{aligned}&-\partial_{\tilde{r}}S_1(\tilde{\theta},\tilde{r})+\partial_{\tilde{r}}S_1(\tilde{\theta}+\tilde{r},\tilde{r})\\&=-\partial_\theta S_1(\tilde{\theta}+\tilde{r},\tilde{r})-\sum_{0<|k|<d}\frac{i(\mathbb{E}v^k)'(\tilde{r})(1-\mu_k(\tilde{r}))+i\mathbb{E}v^k(\tilde{r})\mu_k'(\tilde{r})}{2\pi k}e^{2\pi ik\tilde{\theta}}.\end{aligned}$$

Recalling definitions (20) of $\hat{A}_1$ and (20) of $\hat{B}_1$, this implies that:

$$\hat{A}_1=\mathbb{E}u(\tilde{\theta},\tilde{r})-\mathbb{E}v(\tilde{\theta},\tilde{r})+\hat{B}_1 \tag{26}$$

$$-\sum_{0<|k|<d}\frac{i(\mathbb{E}v^k)'(\tilde{r})(1-\mu_k(\tilde{r}))+i\mathbb{E}v^k(\tilde{r})\mu_k'(\tilde{r})}{2\pi k}e^{2\pi ik\tilde{\theta}}. \tag{27}$$

Then we use (25) and (24) again, noting that $\mu_k'(\tilde{r})=0$ in both regions $|\tilde{r}-p/q|\geq 3\gamma$ and $|\tilde{r}-p/q|\leq\gamma/2$. Moreover, we note that for $|\tilde{r}-p/q|\leq\gamma/2$.

$$\mathbb{E}v_{p,q}(\tilde{\theta},\tilde{r})=\mathbb{E}v_{p,q}(\tilde{\theta},p/q)+\mathcal{O}(\gamma),$$

$$(\mathbb{E}v^k)'(\tilde{r})=(\mathbb{E}v^k)'(p/q)+\mathcal{O}(\gamma).$$

Define

$$E_1(\tilde{\theta},\tilde{r})=-\sum_{0<|k|<d}\frac{i(\mathbb{E}v^k)'(\tilde{r})}{2\pi k}e^{2\pi ik\tilde{\theta}}. \tag{28}$$

Then the same holds for $\mathbb{E}u(\tilde{\theta},\tilde{r})$ and $\mathbb{E}v(\tilde{\theta},\tilde{r})$: recalling definition (15) of E_3, equation (26) yields:

$$\hat{A}_1=\begin{cases}\mathbb{E}u(\tilde{\theta},\tilde{r})-\mathbb{E}v(\tilde{\theta},\tilde{r})+E_1(\tilde{\theta},\tilde{r}) & \text{if}\quad |\tilde{r}-p/q|\geq 3\gamma,\\ \Delta\mathbb{E}(\tilde{\theta},p/q)+\mathbb{E}v_{p,q}(\tilde{\theta})+E_3(\tilde{\theta})+\mathcal{O}(\varepsilon^{1/6}) & \text{if}\quad |\tilde{r}-p/q|\leq\gamma/2,\end{cases} \tag{29}$$

where $\mathbb{E}u(\tilde{\theta},p/q)-\mathbb{E}v(\tilde{\theta},p/q)=\Delta\mathbb{E}(\tilde{\theta},p/q)$. In conclusion, by (29) and (25) we obtain that the first order terms of (17) coincide with the first order terms of (12) and (14) in each region.

For the ε^2-terms we rename $\hat{B}_2$ in the following way:

$$E_2(\tilde{\theta},\tilde{r}) = \hat{B}_2|_{\{|r-p/q|\geq 3\gamma\}}, \tag{30}$$

$$E_4(\tilde{\theta},\tilde{r}) = \hat{B}_2|_{\{|r-p/q|\leq\gamma/2\}}. \tag{31}$$

Now we shall see that E_2 verifies (13). To avoid long notation, in the following we do not write explicitly that expressions A_i, B_i, $\hat{A}_i$ and $\hat{B}_i$ are restricted to the region $\{|r-p/q|\geq 3\gamma\}$. We note that since in this region we have $\hat{B}_1=0$ by (25), recalling the definition (20) of $\hat{B}_1$ it is clear that $B_1=\partial_\theta S_1(\tilde{\theta}+\tilde{r},\tilde{r})$. Hence, from definition (20) of $\hat{B}_2$ it is straightforward to see that:

$$\hat{B}_2=B_2-\partial_\theta^2 S_1(\tilde{\theta}+\tilde{r},\tilde{r})A_1. \tag{32}$$

Now we recall that $\hat{A}_1 = A_1 + \partial_{\tilde{r}} S_1(\tilde{\theta}+\tilde{r},\tilde{r})$. Then, using (29), for $|\tilde{r}-p/q| \geq 3\varepsilon^{1/6}$ we obtain straightforwardly:

$$A_1 = \mathbb{E}u(\tilde{\theta},\tilde{r}) - \mathbb{E}u(\tilde{\theta},\tilde{r}) + E_1(\tilde{\theta},\tilde{r}) - \partial_{\tilde{r}} S_1(\tilde{\theta}+\tilde{r},\tilde{r}). \tag{33}$$

Using this and the definition (18) of B_2 in expression (32) one obtains:

$$\begin{aligned} \hat{B}_2|_{\{|r-p/q|\geq 3\varepsilon^{1/6}\}} \quad = \quad & -\partial_\theta \mathbb{E}v(\tilde{\theta},\tilde{r})\partial_{\tilde{r}} S_1(\tilde{\theta},\tilde{r}) + \partial_r \mathbb{E}v(\tilde{\theta},\tilde{r})\partial_\theta S_1(\tilde{\theta},\tilde{r}) \\ & -\partial_\theta^2 S_1(\tilde{\theta},\tilde{r})\partial_{\tilde{r}} S_1(\tilde{\theta},\tilde{r}) \\ & -\partial_\theta^2 S_1(\tilde{\theta}+\tilde{r},\tilde{r})[\mathbb{E}u(\tilde{\theta},\tilde{r}) - \mathbb{E}u(\tilde{\theta},\tilde{r}) \\ & +E_1(\tilde{\theta},\tilde{r}) - \partial_{\tilde{r}} S_1(\tilde{\theta}+\tilde{r},\tilde{r})]. \end{aligned} \tag{34}$$

Show that this expression has the claimed average (13). On the one hand, it is clear that $\partial_\theta^2 S_1(\tilde{\theta},\tilde{r})\partial_{\tilde{r}} S_1(\tilde{\theta},\tilde{r})$ and $\partial_\theta^2 S_1(\tilde{\theta}+\tilde{r},\tilde{r})\partial_{\tilde{r}} S_1(\tilde{\theta}+\tilde{r},\tilde{r})$ have the same average, so:

$$\int_0^1 -\partial_\theta^2 S_1(\tilde{\theta},\tilde{r})\partial_{\tilde{r}} S_1(\tilde{\theta},\tilde{r}) + \partial_\theta^2 S_1(\tilde{\theta}+\tilde{r},\tilde{r})\partial_{\tilde{r}} S_1(\tilde{\theta}+\tilde{r},\tilde{r}) d\tilde{\theta} = 0.$$

On the other hand, writing explicitly the zeroth Fourier coefficient of the product, one can see that:

$$\int_0^1 -\partial_\theta \mathbb{E}v(\tilde{\theta},\tilde{r})\partial_{\tilde{r}} S_1(\tilde{\theta},\tilde{r}) - \partial_\theta^2 S_1(\tilde{\theta}+\tilde{r},\tilde{r}) E_1(\tilde{\theta},\tilde{r}) d\tilde{\theta} = 0.$$

Thus, recalling (30) and using these two facts in equation (34) we obtain:

$$\int_0^1 E_2(\tilde{\theta},\tilde{r}) d\tilde{\theta} =$$

$$\int_0^1 \left[\partial_r \mathbb{E}v(\tilde{\theta},\tilde{r})\partial_\theta S_1(\tilde{\theta},\tilde{r}) - \partial_\theta^2 S_1(\tilde{\theta},\tilde{r}) \left(\mathbb{E}u(\tilde{\theta},\tilde{r}) - \mathbb{E}v(\tilde{\theta},\tilde{r})\right)\right] d\tilde{\theta},$$

so that (13) is proved.

We note that, from the definition (23) of the Fourier coefficients of S_1, it is clear that S_1 is $\mathcal{C}^l$ with respect to r. Since it just has a finite number of nonzero coefficients, it is analytic with respect to θ. Then, from the definitions (30) of E_2 and (31) of E_4 and the expression (20) of $\hat{B}_2$, it is clear that both E_2 and E_4 are $\mathcal{C}^{l-1}$. Finally we shall bound the $\mathcal{C}^0$-norms of the functions E_2, $b(r)$ and E_4 and also the error terms. To that aim, first let us bound the $\mathcal{C}^l$ norms of S_1 and its derivatives. We will use Lemma 4.1 and proceed similarly as in [5]. We note that:

1. If $\mu_k(\tilde{r}) \neq 1$ we have $|1 - e^{2\pi i k\tilde{r}}| > M\gamma|k|$, and thus:

$$\left|\frac{1}{1 - e^{2\pi i k\tilde{r}}}\right| < M^{-1}\gamma^{-1}|k|^{-1}.$$

2. Then, using that $\|f \circ g\|_{\mathcal{C}^l} \leq C \|f_{|_{\mathrm{Im}(g)}}\|_{\mathcal{C}^l} \left(1 + \|g\|^l_{\mathcal{C}^l}\right)$, we get that:

   $$\left\| \frac{1}{1 - e^{2\pi i k \tilde{r}}} \right\|_{\mathcal{C}^l} \leq M \gamma^{-(l+1)/5} |k|^{-l-1},$$

   for some constant M.

3. Using the rule for the norm of the composition again and the fact that $\|\mu\|_{\mathcal{C}^l}$ is bounded independently of ε, we get:

   $$\|\mu_k(\tilde{r})\|_{\mathcal{C}^l} \leq M \gamma^{-l/5} |k|^{-l},$$

   for some constant M, and the same bound is obtained for $\|1 - \mu_k(\tilde{r})\|_{\mathcal{C}^l}$.

Using items 2 and 3 above and the fact that $\|\mathbb{E}v^k\|_{\mathcal{C}^l}$ are bounded, we get that:

$$\begin{aligned}
\left\| \partial_{\tilde{r}^\alpha} \left[\frac{1 - \mu_k(\tilde{r}) i \mathbb{E}v^k(\tilde{r})}{2\pi k (1 - e^{2\pi i k \tilde{r}})} \right] \right\|_{\mathcal{C}^0} &\leq M_1 \sum_{\alpha_1 + \alpha_2 = \alpha} \frac{1}{2\pi |k|} \|1 - \mu_k(\tilde{r})\|_{\mathcal{C}^{\alpha_1}} \left\| \frac{1}{1 - e^{2\pi i k \tilde{r}}} \right\|_{\mathcal{C}^{\alpha_2}} \\
&\leq M_2 \gamma^{-(\alpha+1)/5} |k|^{-\alpha-2}.
\end{aligned}$$

Then, by item 2 of Lemma 4.1, we obtain:

$$\|S_1\|_{\mathcal{C}^l} \leq M \gamma^{-(l+1)/6}.$$

One can also see that $\|\partial_{\tilde{r}} S_1\|_{\mathcal{C}^l} \leq M \|S_1\|_{\mathcal{C}^{l+1}}$ and $\|\partial_\theta S_1\|_{\mathcal{C}^l} \leq M \|S_1\|_{\mathcal{C}^l}$. In general, one has:

$$\|\partial^n_\theta \partial^m_{\tilde{r}} S_1\|_{\mathcal{C}^l} \leq M \gamma^{-(l+m+1)/6}. \tag{35}$$

Now, recalling definitions (30) of E_2 and (31) of E_4, and using either expression (34) or simply (20), bound (35) yields that there exists some K such that:

$$\|E_2\|_{\mathcal{C}^0} \leq K \gamma^{-1/2}, \qquad \|E_4\|_{\mathcal{C}^0} \leq K \gamma^{-1/2}.$$

To bound $b(r)$ we use again (35) and recall that $\|\mathbb{E}v\|_{\mathcal{C}^0} \leq K$, $\|\partial_r \mathbb{E}v\|_{\mathcal{C}^0} \leq K$, $\|\mathbb{E}u\|_{\mathcal{C}^0} \leq K$. Then from its definition (13) it is clear that:

$$\|b\|_{\mathcal{C}^0} \leq K \gamma.$$

Similarly, one can easily bound the error terms in the equation for $\tilde{r}$:

$$\varepsilon^3 \hat{B}_3 = \mathcal{O}(\varepsilon^{2+a}), \tag{36}$$

and the error terms for the equation of $\tilde{\theta}$:

$$\varepsilon^2 \hat{A}_2 = \mathcal{O}(\varepsilon^{4/3}). \tag{37}$$

This finishes the proof for the normal forms (12) and (14) (in the latter case, we have to take into account the extra error term of order $\mathcal{O}(\varepsilon^{1+a})$ caused by the γ –error term in (29)).

To prove (15), we just need to recall (16) and use (35). Then one obtains:

$$\|\Phi - \mathrm{Id}\|_{\mathcal{C}^2} \leq M'\varepsilon\|S_1\|_{\mathcal{C}^3}.$$

□

From now on we will consider that our deterministic system is in the normal form, and drop tildes.

5 Analysis of the Martingale problem in the strips of each type

5.1 The Totally Irrational case

First of all we note that in this case, as in the IR case, after performing the change to normal form, the n-th iteration of our map can be written as:

$$\begin{aligned} \theta_n &= \theta_0 + nr_0 + \mathcal{O}(n\varepsilon), \\ r_n &= r_0 + \varepsilon \sum_{k=0}^{n-1} \omega_k[v(\theta_k, r_k) + \varepsilon v_2(\theta_k, r_k)] + \varepsilon^2 \sum_{k=0}^{n-1} E_2(\theta_k, r_k) + \mathcal{O}(n\varepsilon^{1+a}), \end{aligned} \tag{38}$$

where $v_2(\theta, r)$ is a given function which can be written explicitly in terms of $v(\theta, r)$ and $S_1(\theta, r)$.

Recall that I_β is a totally irrational segment if $p/q \in I_\beta$, then $|q| > \varepsilon^{-b}$, where $0 < 2b < \beta$.

We recall that we define $b = (\beta - \rho)/2$ for a certain $0 < \rho < \beta$. In the following we shall assume that ρ satisfies an extra condition, which will ensure that certain inequalities are satisfied. This inequalities involve the degree of differentiability of certain $\mathcal{C}^l$ functions. We assume that $l \geq 12$. Then we have that: $\frac{l-11}{l-1} > 0$, $\lim_{l\to\infty} \frac{l-11}{l-1} = 1$. Thus, there exists a constant $R > 0$ such that:

$$\frac{l-11}{l-1} > R > 0, \qquad \text{for all } l \geq 12. \tag{39}$$

Given β, satisfying:

$$0 < \beta \leq 1/5, \tag{40}$$

then we will take ρ satisfying:

$$0 < \rho < R\beta. \tag{41}$$

Lemma 5.1. *Let g be a $\mathcal{C}^l$ function, $l \geq 12$. Suppose r^* satisfies the following condition if for some rational p/q we have $|r^* - p/q| < \varepsilon^\beta$, then $|q| > \varepsilon^{-b}$. Then for any A such that*

$$2\beta < A < (l-1)b - \beta, \quad 0 < \tau = 1 - 2A \leq \min\{A - 2\beta, (l-1)b - A - \beta, \beta\}$$

and ε small enough there is $N \leq \varepsilon^{-A}$ such that for some K independent of ε and any θ^* we have:

$$\left| N \int_0^1 g(\theta, r^*) d\theta - \sum_{k=0}^{N-1} g(\theta^* + kr^*, r^*) \right| \leq K\varepsilon^{\tau+\beta}.$$

In particular, one can choose any $0 < \beta \leq 1/5$, $A = 7\beta/3$, $\tau = A - 2\beta = \beta/3$, $b = \beta/3$.

Proof. Denote $g_0(r) = \int_0^1 g(\theta, r)d\theta$. Expand $g(\theta, r)$ in its Fourier series, i.e.:

$$g(\theta, r) = g_0(r) + \sum_{m \in \mathbb{Z}\setminus\{0\}} g_m(r) e^{2\pi i m\theta}$$

for some $g_m(r) : \mathbb{R} \to \mathbb{C}$. Then we have:

$$\begin{aligned}
&\sum_{k=0}^{N-1} (g(\theta^* + kr^*, r^*) - g_0(r^*)) = \sum_{k=0}^{N-1} \sum_{m \in \mathbb{Z}\setminus\{0\}} g_m(r^*) e^{2\pi i m(\theta^* + kr^*, r^*))} \\
&= \sum_{k=0}^{N-1} \sum_{1 \leq |m| \leq [\varepsilon^{-b}]} g_m(r^*) e^{2\pi i m(\theta^* + kr^*)} + \sum_{k=0}^{N} \sum_{|m| \geq [\varepsilon^{-b}]} g_m(r^*) e^{2\pi i m(\theta^* + kr^*)} \\
&= \sum_{1 \leq |m| \leq [\varepsilon^{-b}]} g_m(r^*) e^{2\pi i m\theta^*} \sum_{k=0}^{N-1} e^{2\pi i m k r^*} + \sum_{k=0}^{N-1} \sum_{|m| \geq [\varepsilon^{-b}]} g_m(r^*) e^{2\pi i m(\theta^* + kr^*)} \\
&= \sum_{1 \leq |m| \leq [\varepsilon^{-b}]} g_m(r^*) e^{2\pi i m\theta^*} \frac{e^{2\pi i N m r^*} - 1}{e^{2\pi i m r^*} - 1} + \sum_{k=0}^{N-1} \sum_{|m| \geq [\varepsilon^{-b}]} g_m(r^*) e^{2\pi i m(\theta^* + kr^*)}.
\end{aligned} \tag{42}$$

To bound the first sum in (42) we distinguish into the following cases:

- If r^* is rational p/q, we know that $|q| > \varepsilon^{-b}$.
  - If $|q| \leq \varepsilon^{-A}$, then pick $N = |q|$ and the first sum vanishes.
  - If $|q| > \varepsilon^{-A}$, then by definition of r^* for any s/m with $|m| < \varepsilon^{-b}$ we have or $|mr^* - s| > \varepsilon^\beta$. By the pigeon hole principle there exist integers $0 < N = \tilde{q} < \varepsilon^{-A}$ and $\tilde{p}$ such that $|\tilde{q}r^* - \tilde{p}| \leq 2\varepsilon^A$.

- If r^* is irrational, consider a continuous fraction expansion $p_n/q_n \to r^*$ as $n \to \infty$. Choose $p'/q' = p_n/q_n$ with n such that $q_{n+1} > \varepsilon^{-A}$. This implies that $|q'r^* - p'| < 1/|q_{n+1}| \leq \varepsilon^A$. The same argument as above shows that for any $|m| < \varepsilon^{-b}$ we have $|mr^* - s| > \varepsilon^\beta$.

Let N be as above. Then

$$\left| \sum_{1 \leq |m| \leq [\varepsilon^{-b}]} g_m(r^*)\; e^{2\pi i m\theta^*} \frac{e^{2\pi i N m r^*} - 1}{e^{2\pi i m r^*} - 1} \right| \leq 2\varepsilon^{A-\beta} \sum_{1 \leq |m| \leq [\varepsilon^{-b}]} |g_m(r^*)|.$$

We point out that since $g(\theta, r)$ is $\mathcal{C}^l$, then its Fourier coefficients satisfy $|g_m(r^*)| \leq C|m|^{-l},\ m \neq 0$. Thus we can bound the first sum in (42) by:

$$\left| \sum_{1 \leq |m| \leq [\varepsilon^{-b}]} g_m(r^*)\; e^{2\pi i m\theta^*} \frac{e^{2\pi i N m r^*} - 1}{e^{2\pi i m r^*} - 1} \right| \tag{43}$$

$$\leq \varepsilon^{A-\beta} \sum_{1 \leq |m| \leq [\varepsilon^{-b}]} |g_m(r^*)| \;\leq\; C\varepsilon^{A-\beta} \sum_{1 \leq |m| \leq [\varepsilon^{-b}]} \frac{1}{m^2} \leq K\varepsilon^{A-\beta}. \tag{44}$$

To bound the second sum we use again the bound for the Fourier coefficients $g_m(r^*)$:

$$\left| \sum_{k=0}^{N} \sum_{|m| \geq [\varepsilon^{-b}]} g_m(r^*) e^{2\pi i m(\theta + k r^*)} \right| \leq N \sum_{|m| \geq [\varepsilon^{-b}]} \frac{1}{m^l} \leq \tag{45}$$

$$KN\varepsilon^{(l-1)b} \leq K\varepsilon^{(l-1)b-A}.$$

Clearly, taking $\tau = 1 - 2A \leq \min\{A - 2\beta, (l-1)b - A - \beta, \beta\}$, and substituting (43) and (45) in (42) we obtain the claim of the lemma. □

Lemma 5.2. *Let β satisfy (40), and $b = (\beta - \rho)/2$ with ρ satisfying (41). Let n_β be an exit time of the process (θ_n, r_n) defined by (38) from some bounded domain I_β. Let $\delta > 0$ be small enough. Suppose that $n_\beta \geq \varepsilon^{-2(1-\beta)+\delta}$. For all $l \geq 8$ the following holds:*

1. *Given two $\mathcal{C}^l$ functions $h : \mathbb{R} \to \mathbb{R}$ and $g : \mathbb{T} \times \mathbb{R} \to \mathbb{R}$, there exists a constant $d > 0$ such that:*

$$\varepsilon^2 \sum_{k=0}^{n_\beta - 1} e^{-\lambda \varepsilon^2 k} h(r_k)(g(\theta_k, r_k) - g_0(r_k)) = \mathcal{O}(\varepsilon^d),$$

   where $g_0(r) = \int_0^1 g(\theta, r) d\theta$.

2. *If $n_\beta < \varepsilon^{-2(1-\beta)+\delta}$, then given a $\mathcal{C}^l$ function $g : \mathbb{T} \times \mathbb{R} \to \mathbb{R}$ and a collection of functions $h_k : \mathbb{R} \to \mathbb{R}$, with $\|h_k\|_{\mathcal{C}^0} \le M$ and $\|h_{k+1} - h_k\|_{\mathcal{C}^0} \le M\varepsilon^2$ for all k, there exists a constant $d > 0$ such that*

$$\varepsilon^2 \sum_{k=0}^{n_\beta - 1} h_k(r_k)(g(\theta_k, r_k) - g_0(r_k)) = \mathcal{O}(\varepsilon^{2\beta+d}),$$

where $g_0(r) = \int_0^1 g(\theta, r) d\theta$.

Proof. We shall prove both claims using Lemma 5.1. To that aim fix $2\beta < A < \min\{(l-1)b - \beta, (1-\beta)/2\}$. We note that (40) and (41) ensure that $2\beta < \min\{(l-1)b - \beta, (1-\beta)/2\}$, so there always exists such A.

Since we have $n_\beta \ge \varepsilon^{-(1-\beta)+\delta}$ and we have taken $A < (1-\beta)/2 < 1-\beta-\delta$ (the second inequality being satisfied because δ is small), we have $\varepsilon^{-A} < n_\beta$. Choose $N < \varepsilon^{-A}$ from Lemma 5.1 and write $n_\beta = P_\beta N + Q_\beta$, for some integers P_β and $0 \le Q_\beta < N$. Then:

$$\begin{aligned} & \varepsilon^2 \left| \sum_{k=0}^{n_\beta-1} e^{-\lambda\varepsilon^2 k} h(r_k)(g(\theta_k, r_k) - g_0(r_k)) \right| \\ \le \quad & \varepsilon^2 \left| \sum_{k=0}^{P_\beta - 1} \sum_{j=0}^{N-1} e^{-\lambda\varepsilon^2(kN+j)} h(r_{kN+j})(g(\theta_{kN+j}, r_{kN+j}) - g_0(r_{kN+j})) \right| \qquad (46) \\ + \quad & \varepsilon^2 \left| \sum_{j=0}^{Q_\beta - 1} e^{-\lambda\varepsilon^2(P_\beta N+j)} h(r_{P_\beta N+j})(g(\theta_{P_\beta N+j}, r_{P_\beta N+j}) - g_0(r_{P_\beta N+j})) \right|. \end{aligned}$$

Let us prove item 1. We shall bound the two terms in the right hand side of (46) in a different way. Recall that in the normal form (12) we have

$$f_\omega \begin{pmatrix} \theta \\ r \end{pmatrix} \mapsto \begin{pmatrix} \theta + r + \varepsilon \mathbb{E} u(\theta, r) + \varepsilon \omega u(\theta, r) + \mathcal{O}_l(\varepsilon^{1+a}) \\ r + \varepsilon^2 E_2(\theta, r) + \mathcal{O}_l(\varepsilon^{2+a}). \end{pmatrix}. \qquad (47)$$

On the one hand we have that for all $k \le P_\beta$, and all $j \le N$:

$$r_{kN+j} = r_{kN} + \mathcal{O}(N\varepsilon^2), \qquad (48)$$

and:

$$\theta_{kN+j} = \theta_{kN} + j r_{kN} + \mathcal{O}(N^2\varepsilon). \qquad (49)$$

Hence:

$$\begin{aligned} & e^{-\lambda\varepsilon^2(kN+j)} h(r_{kN+j})(g(\theta_{kN+j}, r_{kN+j}) - g_0(r_{kN+j})) \\ = \quad & e^{-\lambda\varepsilon^2 kN} h(r_{kN})(g(\theta_{kN} + j r_{kN}, r_{kN}) - g_0(r_{kN})) + \mathcal{O}(e^{-\lambda\varepsilon^2 kN} N^2 \varepsilon). \end{aligned}$$

Then:

$$\begin{aligned}
&\varepsilon^2\left|\sum_{k=0}^{P_\beta-1}\sum_{j=0}^{N-1}e^{-\lambda\varepsilon^2(kN+j)}h(r_{kN+j})(g(\theta_{kN+j},r_{kN+j})-g_0(r_{kN+j}))\right| \\
\leq\quad &\varepsilon^2\left|\sum_{k=0}^{P_\beta-1}e^{-\lambda\varepsilon^2kN}h(r_{kN})\sum_{j=0}^{N-1}(g(\theta_{kN}+jr_{kN},r_{kN})-g_0(r_{kN}))\right| \\
&+KN^3\varepsilon^3\sum_{k=0}^{P_\beta-1}e^{-\lambda\varepsilon^2kN}.
\end{aligned}$$

Thus, using Lemma 5.1 we obtain:

$$\begin{aligned}
&\varepsilon^2\left|\sum_{k=0}^{P_\beta-1}e^{-\lambda\varepsilon^2kN}h(r_{kN})\sum_{j=0}^{N-1}(g(\theta_{kN}+jr_{kN},r_{kN})-g_0(r_{kN}))\right| \\
\leq\quad &K\varepsilon^{2+\tau+\beta}\sum_{k=0}^{P_\beta-1}e^{-\lambda\varepsilon^2kN}|h(r_{kN})|\leq\tilde{K}\varepsilon^{\tau+\beta},
\end{aligned}$$

for some constants $K,\tilde{K}>0$. Moreover, we have:

$$KN^3\varepsilon^3\sum_{k=0}^{P_\beta-1}e^{-\lambda\varepsilon^2kN}\leq\tilde{K}\varepsilon^{1+\beta-2A}.$$

Thus:

$$\begin{aligned}
&\varepsilon^2\left|\sum_{k=0}^{P_\beta-1}\sum_{j=0}^{N-1}e^{-\lambda\varepsilon^2(kN+j)}h(r_{kN+j})(g(\theta_{kN+j},r_{kN+j})-g_0(r_{kN+j}))\right| \\
&\qquad\leq K(\varepsilon^{\tau+\beta}+\varepsilon^{1+\beta-2A}).
\end{aligned}\tag{50}$$

On the other hand we have:

$$\begin{aligned}
&\varepsilon^2\left|\sum_{j=0}^{Q_\beta-1}e^{-\lambda\varepsilon^2(P_\beta N+j)}h(r_{P_\beta N+j})(g(\theta_{P_\beta N+j},r_{P_\beta N+j})-g_0(r_{P_\beta N+j}))\right| \\
&\leq\varepsilon^2K\sup_{(\theta,r)\in I_\beta}|h(r)(g(\theta,r)-g_0(r))|Q_\beta\leq\tilde{K}\varepsilon^{2-A}.
\end{aligned}\tag{51}$$

In conclusion, using (50) and (51) in equation (46) we obtain:

$$\varepsilon^2\left|\sum_{k=0}^{n_\beta-1}e^{-\lambda\varepsilon^2k}h(r_k)(g(\theta_k,r_k)-g_0(r_k))\right|\leq K(\varepsilon^{2-A}+\varepsilon^{1+\beta-2A}+\varepsilon^{\tau+\beta}).$$

Denoting $d = \min\{2-A, 1+\beta-2A, \tau+\beta\}$, and letting $1-2A = \tau \geq \beta/3$. the proof of item 1 is finished.

Now let us prove item 2. The proof is very similar to item 1. We can use the same formula (46) substituting $e^{-\lambda\varepsilon^2 k} h(r_k)$ by $h_k(r_k)$. Again, using (48) and (49) we can write:

$$\begin{aligned} & h_{kN+j}(r_{kN+j})(g(\theta_{kN+j}, r_{kN+j}) - g_0(r_{kN+j})) \\ = \quad & h_{kN}(r_{kN})(g(\theta_{kN} + jr_{kN}, r_{kN}) - g_0(r_{kN})) + \mathcal{O}(N^2\varepsilon). \end{aligned}$$

Then:

$$\begin{aligned} & \varepsilon^2 \left| \sum_{k=0}^{P_\beta-1} \sum_{j=0}^{N-1} h_{kN+j}(r_{kN+j})(g(\theta_{kN+j}, r_{kN+j}) - g_0(r_{kN+j})) \right| \\ \leq \quad & \varepsilon^2 \left| \sum_{k=0}^{P_\beta-1} h_{kN}(r_{kN}) \sum_{j=0}^{N-1} (g(\theta_{kN} + jr_{kN}, r_{kN}) - g_0(r_{kN})) \right| + KN^3\varepsilon^3 P_\beta. \end{aligned}$$

Thus, using Lemma 5.1 we obtain:

$$\varepsilon^2 \left| \sum_{k=0}^{P_\beta-1} h_{kN}(r_{kN}) \sum_{j=0}^{N-1} (g(\theta_{kN} + jr_{kN}, r_{kN}) - g_0(r_{kN})) \right| \leq$$

$$K\varepsilon^{2+\tau+\beta} P_\beta \leq K\varepsilon^{\tau+2\beta},$$

where we have used that $P_\beta \leq n_\beta \leq \varepsilon^{-2+2\beta-\delta} \leq \varepsilon^{-2+\beta}$. For the same reason we have $K\varepsilon^{3-2A} P_\beta \leq K\varepsilon^{1-2A+\beta}$.Thus:

$$\begin{aligned} & \varepsilon^2 \left| \sum_{k=0}^{P_\beta-1} \sum_{j=0}^{N-1} h_{kN+j}(r_{kN+j})(g(\theta_{kN+j}, r_{kN+j}) - g_0(r_{kN+j})) \right| \leq \\ & K(\varepsilon^{\tau+2\beta} + \varepsilon^{1-2A+\beta}). \end{aligned} \tag{52}$$

On the other hand we have:

$$\begin{aligned} & \varepsilon^2 \left| \sum_{j=0}^{Q_\beta-1} h_{P_\beta N+j}(r_{P_\beta N+j})(g(\theta_{P_\beta N+j}, r_{P_\beta N+j}) - g_0(r_{P_\beta N+j})) \right| \\ & \leq \varepsilon^2 K Q_\beta \leq \tilde{K}\varepsilon^{2-A}. \end{aligned} \tag{53}$$

In conclusion, using (52) and (53) in equation (46) we obtain:

$$\varepsilon^2 \left| \sum_{k=0}^{n_\beta-1} e^{-\lambda\varepsilon^2 k} h_k(r_k)(g(\theta_k, r_k) - g_0(r_k)) \right| \leq K(\varepsilon^{2-A} + \varepsilon^{1-2A+\beta} + \varepsilon^{\tau+2\beta}).$$

Choosing $\tau = 1-2A > 0$ the last two terms are the same. In particular, $A < 1/2$ and $2-A > 3/2$. Therefore, the first term is negligible. $\square$

Let r_0 belong to the TI case. Consider an interval $I_\beta = \{(\theta, r) \in \mathbb{T} \times \mathbb{R} : |r - r_0| \le \varepsilon^\beta\}$, for some $0 < \beta \le 1/5$. Denote $n_\beta \in \mathbb{N}$ the exit time from I_β, that is the first number such that $(\theta_{n_\beta}, r_{n_\beta}) \notin I_\beta$.

Lemma 5.3. *Let β satisfy* (40)*, and $b = (\beta - \rho)/2$ with ρ satisfying* (41)*. Take $f : \mathbb{R} \to \mathbb{R}$ be any $\mathcal{C}^l$ function with $l \ge 12$. Then there exists $d > 0$ such that for all $\lambda > 0$ one has:*

$$\mathbb{E}\Big(e^{-\lambda\varepsilon^2 n_\beta} f(r_{n_\beta}) + \varepsilon^2 \sum_{k=0}^{n_\beta - 1} e^{-\lambda\varepsilon^2 k}\Big[\lambda f(r_k) - \Big(b(r_k) f'(r_k) + \frac{\sigma^2(r_k)}{2} f''(r_k)\Big)\Big]\Big) - f(r_0) = \mathcal{O}(\varepsilon^{2\beta + d}),$$

where for $E_2(\theta, r)$, defined in (11), we have

$$b(r) = \int_0^1 E_2(\theta, r)\, d\theta, \qquad \sigma^2(r) = \int_0^1 v^2(\theta, r)\, d\theta.$$

Proof. Let us denote:

$$\eta = e^{-\lambda\varepsilon^2 n_\beta} f(r_{n_\beta}) + \varepsilon^2 \sum_{k=0}^{n_\beta - 1} e^{-\lambda\varepsilon^2 k}\Big[\lambda f(r_k) - \Big(b(r_k) f'(r_k) + \frac{\sigma^2(r_k)}{2} f''(r_k)\Big)\Big]. \tag{54}$$

First of all we shall use the law of total expectation. Fix a small enough $\delta > 0$. Then we have:

$$\begin{aligned}\mathbb{E}(\eta) &= \mathbb{E}\big(\eta \,|\, \varepsilon^{-2(1-\beta)+\delta} \le n_\beta \le \varepsilon^{-2(1-\beta)-\delta}\big)\, \mathbb{P}\{\varepsilon^{-2(1-\beta)+\delta} \le n_\beta \le \varepsilon^{-2(1-\beta)-\delta}\} \\ &+ \mathbb{E}\big(\eta \,|\, n_\beta < \varepsilon^{-2(1-\beta)+\delta}\big)\, \mathbb{P}\{n_\beta < \varepsilon^{-2(1-\beta)+\delta}\} \\ &+ \mathbb{E}\big(\eta \,|\, n_\beta > \varepsilon^{-2(1-\beta)-\delta}\big)\, \mathbb{P}\{n_\beta > \varepsilon^{-2(1-\beta)-\delta}\}.\end{aligned}$$

By Lemma B.2 for ε sufficiently small and $c > 0$ independent of ε we have

$$\mathbb{P}\{n_\beta < \varepsilon^{-2(1-\beta)+\delta}\} \le \exp\Big(-\frac{c}{\varepsilon^{2\delta}}\Big). \tag{55}$$

Now we write:

$$e^{-\lambda\varepsilon^2 n_\beta} f(r_{n_\beta}) = f(r_0) + \sum_{k=0}^{n_\beta - 1}\Big(e^{-\lambda\varepsilon^2(k+1)} f(r_{k+1}) - e^{-\lambda\varepsilon^2 k} f(r_k)\Big).$$

Doing the Taylor expansion in each term inside the sum we get:

$$\begin{aligned}e^{-\lambda\varepsilon^2 n_\beta} f(r_{n_\beta}) &= f(r_0) + \sum_{k=0}^{n_\beta - 1}\Big[-\lambda\varepsilon^2 e^{-\lambda\varepsilon^2 k} f(r_k) + e^{-\lambda\varepsilon^2 k} f'(r_k)(r_{k+1} - r_k) \\ &\quad + \frac{1}{2} e^{-\lambda\varepsilon^2 k} f''(r_k)(r_{k+1} - r_k)^2 + \mathcal{O}(e^{-\lambda\varepsilon^2 k}\varepsilon^3)\Big].\end{aligned}$$

Substituting this in (54) we get:

$$\begin{aligned}\eta &= f(r_0)+\sum_{k=0}^{n_\beta-1}\left[e^{-\lambda\varepsilon^2 k}f'(r_k)(r_{k+1}-r_k)+\frac{1}{2}e^{-\lambda\varepsilon^2 k}f''(r_k)(r_{k+1}-r_k)^2\right]\\ &\quad -\varepsilon^2\sum_{k=0}^{n_\beta-1}e^{-\lambda\varepsilon^2 k}\left[b(r_k)f'(r_k)+\frac{\sigma^2(r_k)}{2}f''(r_k)\right]+\sum_{k=0}^{n_\beta-1}\mathcal{O}(e^{-\lambda\varepsilon^2 k}\varepsilon^3). \quad (56)\end{aligned}$$

We note that using (38) we can write:

$$r_{k+1}-r_k=\varepsilon\omega_k[v(\theta_k,r_k)+\varepsilon v_2(\theta_k,r_k)]+\varepsilon^2E_2(\theta_k,r_k)+\mathcal{O}(\varepsilon^{2+a}),$$

and also:

$$(r_{k+1}-r_k)^2=\varepsilon^2v^2(\theta_k,r_k)+\mathcal{O}(\varepsilon^3).$$

Thus we can rewrite (56) as:

$$\begin{aligned}\eta &= f(r_0)+\varepsilon\sum_{k=0}^{n_\beta-1}e^{-\lambda\varepsilon^2 k}f'(r_k)\omega_k\left[v(\theta_k,r_k)+\varepsilon v_2(\theta_k,r_k)\right]\\ &\quad +\varepsilon^2\sum_{k=0}^{n_\beta-1}e^{-\lambda\varepsilon^2 k}f'(r_k)\left[E_2(\theta_k,r_k)-b(r_k)\right]\\ &\quad +\frac{\varepsilon^2}{2}\sum_{k=0}^{n_\beta-1}e^{-\lambda\varepsilon^2 k}f''(r_k)\left[v^2(\theta_k,r_k)-\sigma^2(r_k)\right]\\ &\quad +\sum_{k=0}^{n_\beta-1}\mathcal{O}(e^{-\lambda\varepsilon^2 k}\varepsilon^{2+a}). \quad (57)\end{aligned}$$

Now we distinguish between the case $\varepsilon^{-2(1-\beta)+\delta}\le n_\beta\le\varepsilon^{-2(1-\beta)-\delta}$ and $n_\beta>\varepsilon^{-2(1-\beta)-\delta}$. Consider the former case. First, we show that the last term in (57) is $\mathcal{O}(\varepsilon^{\beta+d})$ for some $d>0$. Indeed,

$$\left|\sum_{k=0}^{n_\beta-1}\mathcal{O}(e^{-\lambda\varepsilon^2 k}\varepsilon^{2+a})\right|\le K\varepsilon^{2+a}n_\beta\le K\varepsilon^{2\beta+d}, \quad (58)$$

where $d=a-\delta>0$ due to smallness of δ, and K is some positive constant. Now we use item 2 of Lemma 5.2 in (57) twice. First we take $h_k(r)=e^{-\lambda\varepsilon^2 k}f'(r)$ and $g(\theta,r)=E_2(\theta,r)$, and after we take $h_k(r)=e^{-\lambda\varepsilon^2 k}f''(r)$ and $g(\theta,r)=v^2(\theta,r)$. Then, recalling also (58), equation (57) for $\varepsilon^{-2(1-\beta)+\delta}\le n_\beta\le\varepsilon^{-2(1-\beta)-\delta}$ yields:

$$\eta=f(r_0)+\varepsilon\sum_{k=0}^{n_\beta-1}e^{-\lambda\varepsilon^2 k}f'(r_k)\omega_k\left[v(\theta_k,r_k)+\varepsilon v_2(\theta_k,r_k)\right]+\mathcal{O}(\varepsilon^{2\beta+d}). \quad (59)$$

Now we focus on the case $n_\beta > \varepsilon^{-2(1-\beta)-\delta}$. The last term in (57) can be bounded by:

$$\left|\sum_{k=0}^{n_\beta-1} \mathcal{O}(e^{-\lambda\varepsilon^2 k}\varepsilon^{2+a})\right| \leq K\varepsilon^{2+a}\sum_{k=0}^{n_\beta-1} e^{-\lambda\varepsilon^2 k} = K\varepsilon^{2+a}\frac{1-e^{-\lambda\varepsilon^2 n_\beta}}{1-e^{-\lambda\varepsilon^2}} \leq K_\lambda\varepsilon^a, \tag{60}$$

for some positive constants K and K_λ. Similarly to the previous case, we use item 1 of Lemma 5.2 in (57) twice. First we take $h(r) = f'(r)$ and $g(\theta, r) = E_2(\theta, r)$, and after we take $h(r) = f''(r)$ and $g(\theta, r) = v^2(\theta, r)$. Using this and bound (60) in equation (57), we obtain the following bound for $n_\beta > \varepsilon^{-2(1-\beta)-\delta}$:

$$\eta = f(r_0) + \varepsilon\sum_{k=0}^{n_\beta-1} e^{-\lambda\varepsilon^2 k} f'(r_k)\omega_k\left[v(\theta_k, r_k) + \varepsilon v_2(\theta_k, r_k)\right] + \mathcal{O}(\varepsilon^d). \tag{61}$$

Now we just need to note that since ω_k is independent of r_k and θ_k, we have for all $k \in \mathbb{N}$:

$$\begin{aligned} \mathbb{E}(\omega_k f'(r_k)[v(\theta_k, r_k) + \varepsilon v_2(\theta_k, r_k)]) &= \\ \mathbb{E}(\omega_k)\mathbb{E}(f'(r_k)[v(\theta_k, r_k) + \varepsilon v_2(\theta_k, r_k)]) &= 0, \end{aligned}$$

because $\mathbb{E}(\omega_k) = 0$. Thus, denoting $n_\varepsilon = [\varepsilon^{-2(1-\beta)+\delta}]$, if we take expectations in (59) and (61) and use the total expectation formula, it is clear that:

$$\begin{aligned} & \mathbb{E}(\eta) - f(r_0) \\ = \;& \varepsilon \sum_{\substack{n\in\mathbb{N} \\ n\geq n_\varepsilon}} \mathbb{E}\left(\sum_{k=0}^{n-1} e^{-\lambda\varepsilon^2 k} f'(r_k)\omega_k\left[v(\theta_k, r_k) + \varepsilon v_2(\theta_k, r_k]\right)\right)\mathbb{P}\{n_\beta = n\} \\ + \;& \mathcal{O}(\varepsilon^{2\beta+d})\mathbb{P}\{\varepsilon^{-2(1-\beta)+\delta} \leq n_\beta \leq \varepsilon^{-2(1-\beta)-\delta}\} + \mathcal{O}(\varepsilon^d)\mathbb{P}\{n_\beta > \varepsilon^{-2(1-\beta)-\delta}\} \\ = \;& \varepsilon \sum_{\substack{n\in\mathbb{N} \\ n\geq n_\varepsilon}} \left(\sum_{k=0}^{n-1} e^{-\lambda\varepsilon^2 k}\mathbb{E}(f'(r_k)\omega_k\left[v(\theta_k, r_k) + \varepsilon v_2(\theta_k, r_k)\right])\right)\mathbb{P}\{n_\beta = n\} \\ + \;& \mathcal{O}(\varepsilon^{2\beta+d})\mathbb{P}\{\varepsilon^{-2(1-\beta)+\delta} \leq n_\beta \leq \varepsilon^{-2(1-\beta)-\delta}\} + \mathcal{O}(\varepsilon^d)\mathbb{P}\{n_\beta > \varepsilon^{-2(1-\beta)-\delta}\} \\ = \;& \mathcal{O}(\varepsilon^{2\beta+d})\mathbb{P}\{\varepsilon^{-2(1-\beta)+\delta} \leq n_\beta \leq \varepsilon^{-2(1-\beta)-\delta}\} + \mathcal{O}(\varepsilon^d)\mathbb{P}\{n_\beta > \varepsilon^{-2(1-\beta)-\delta}\} \end{aligned} \tag{62}$$

By Lemma B.2 there exists a constant $a > 0$ such that:

$$\mathbb{P}\{n_\beta > \varepsilon^{-2(1-\beta)-\delta}\} = \mathcal{O}\left(e^{-\frac{a}{\varepsilon^\delta}}\right). \tag{63}$$

Clearly, if (63) is true then $\mathbb{P}\{n_\beta > \varepsilon^{-2(1-\beta)-\delta}\}$ is smaller than any order of ε and then (62) finishes the proof of the lemma.

To prove (63), let us define $n_\delta := [\varepsilon^{-2(1-\beta)-\delta}]$. Define also $n_i := i[\varepsilon^{-2(1-\beta)}]$, $i = 0, \dots, [\varepsilon^{-\delta}]$. Clearly, if $n_\beta > \varepsilon^{-2(1-\beta)-\delta}$, then $|r_{n_{i+1}} - r_{n_i}| < 2\varepsilon^\beta$ for all i. In other words, we have that:

$$\begin{aligned} \mathbb{P}\{n_\beta > \varepsilon^{-2(1-\beta)-\delta}\} &\leq \mathbb{P}\{|r_{n_{i+1}} - r_{n_i}| < 2\varepsilon^\beta \text{ for all } i = 0, \dots, [\varepsilon^{-\delta}]\} \\ &= \prod_{i=0}^{[\varepsilon^{-\delta}]} \mathbb{P}\{|r_{n_{i+1}} - r_{n_i}| < 2\varepsilon^\beta\}, \end{aligned} \tag{64}$$

where in the last equality we have used that $r_{n_{i+1}} - r_{n_i}$ and $r_{n_{j+1}} - r_{n_j}$ are independent if $i \neq j$.

Now, take any i. Then:

$$r_{n_{i+1}} - r_{n_i} = \varepsilon \sum_{i=n_i}^{n_{i+1}-1} \omega_k v(\theta_k, r_k) + \mathcal{O}(\varepsilon^2(n_i - n_{i+1})).$$

Note that $\varepsilon^2(n_i - n_{i+1}) = \varepsilon^2[\varepsilon^{-2(1-\beta)}] \leq \varepsilon^{2\beta}$. Thus:

$$\varepsilon \left| \sum_{i=n_i}^{n_{i+1}-1} \omega_k v(\theta_k, r_k) \right| - \mathcal{O}(\varepsilon^{2\beta}) \leq |r_{n_{i+1}} - r_{n_i}|. \tag{65}$$

As a consequence, if $|r_{n_{i+1}} - r_{n_i}| \leq 2\varepsilon^\beta$ then $\varepsilon \left|\sum_{i=n_i}^{n_{i+1}-1} \omega_k v(\theta_k, r_k)\right| \leq 3\varepsilon^\beta$. Indeed, if this latter inequality does not hold, then:

$$\varepsilon \left| \sum_{i=n_i}^{n_{i+1}-1} \omega_k v(\theta_k, r_k) \right| - \mathcal{O}(\varepsilon^{2\beta}) > 3\varepsilon^\beta(1 - \mathcal{O}(\varepsilon^\beta)) \geq 2\varepsilon^\beta \geq |r_{n_{i+1}} - r_{n_i}|,$$

which is a contraditciton with (65). In other words:

$$\mathbb{P}\{|r_{n_{i+1}} - r_{n_i}| < 2\varepsilon^\beta\} \leq \mathbb{P}\left\{ \varepsilon \left| \sum_{i=n_i}^{n_{i+1}-1} \omega_k v(\theta_k, r_k) \right| \leq 3\varepsilon^\beta \right\}.$$

Now by Lemma B.2 that $(n_{i+1} - n_i)^{-1/2} \sum_{k=n_i}^{n_{i+1}-1} \omega_k v(\theta_k, r_k)$ converges in distribution to $\xi \sim \mathcal{N}(0, c^2)$ for some $c > 0$.

Thus, using that $n_{i+1} - n_i = [\varepsilon^{-2(1-\beta)}]$, as $\varepsilon \to 0$ we obtain:

$$\mathbb{P}\left\{ \varepsilon \left| \sum_{i=n_i}^{n_{i+1}-1} \omega_k v(\theta_k, r_k) \right| \leq 3\varepsilon^\beta \right\} = \mathbb{P}\{|\xi| \leq 3\} + o(1) \leq \rho < 1,$$

for some constant $\rho > 0$. Then, using this in (64) we get:

$$\mathbb{P}\{n_\beta > \varepsilon^{-2(1-\beta)-\delta}\} \leq \rho^{1/\varepsilon^\delta}.$$

Defining $a = -\log \rho > 0$ (because $\rho < 1$) we obtain claim (63). $\square$

5.2 The Imaginary Rational case

In this section we deal with the imaginary rational case. The ideas are basically the same as in the TI case. Recall that after performing the change to normal form the n-th iteration of our map can be written as:

$$\begin{aligned} \theta_n &= \theta_0 + nr_0 + \mathcal{O}(n\varepsilon), \\ r_n &= r_0 + \varepsilon \sum_{k=0}^{n-1} \omega_k [v(\theta_k, r_k) + \varepsilon v_2(\theta_k, r_k)] \\ &\quad + \varepsilon^2 \textstyle\sum_{k=0}^{n-1} E_2(\theta_k, r_k) + \mathcal{O}(n\varepsilon^{2+a}), \end{aligned} \tag{66}$$

where $v_2(\theta, r)$ is a given function which can be written explicitly in terms of $v(\theta, r)$ and $S_1(\theta, r)$.

We also recall that given an imaginary rational strip I_β there exists a unique $r^* \in I_\beta$, with $r^* = p/q$ and $|q| < \varepsilon^{-b}$. Moreover, for all $r_0 \in I_\beta$ we have $|r_0 - r^*| \leq \varepsilon^\beta$. Then by (66) for any $n \leq n_\beta$ we have:

$$\begin{aligned} \theta_n = &\quad \theta_0 + nr^* + \mathcal{O}(n\varepsilon^\beta), \\ r_n = &\qquad r^* + \mathcal{O}(\varepsilon^\beta). \end{aligned} \tag{67}$$

Define

$$g_0(\theta, r) = \frac{1}{q} \sum_{i=0}^{q-1} g(\theta + ir, r). \tag{68}$$

Lemma 5.4. *Let β satisfy (40), and $b = (\beta - \rho)/2$ with ρ satisfying (41). Let n_β be an exit time of the process (θ_n, r_n) defined by (38) from some bounded domain I_β. For all $l \geq 1$ the following holds:*

1. *Given two $\mathcal{C}^l$ functions $h : \mathbb{R} \to \mathbb{R}$ and $g : \mathbb{T} \times \mathbb{R} \to \mathbb{R}$, there exists a constant $d > 0$ such that:*

$$\varepsilon^2 \sum_{k=0}^{n_\beta - 1} e^{-\lambda \varepsilon^2 k} h(r_k)(g(\theta_k, r_k) - g_0(\theta_k, r_k)) = \mathcal{O}(\varepsilon^d).$$

2. *If $n_\beta < \varepsilon^{-2+\beta}$, then given a $\mathcal{C}^l$ function $g : \mathbb{T} \times \mathbb{R} \to \mathbb{R}$ and a collection of functions $h_k : \mathbb{R} \to \mathbb{R}$, with $\|h_k\|_{\mathcal{C}^0} \leq M$ and $\|h_{k+1} - h_k\|_{\mathcal{C}^0} \leq M\varepsilon^2$ for all k, there exists a constant $d > 0$ such that:*

$$\varepsilon^2 \sum_{k=0}^{n_\beta - 1} h_k(r_k)(g(\theta_k, r_k) - g_0(\theta_k, r_k)) = \mathcal{O}(\varepsilon^{2\beta + d}).$$

Proof. Let us start with item 1. Write $n_\beta = P_\beta q + Q_\beta$, for some integers P_β and $0 \leq Q_\beta < q$. Then:

$$\begin{aligned}
& \varepsilon^2 \left| \sum_{k=0}^{n_\beta - 1} e^{-\lambda \varepsilon^2 k} h(r_k)(g(\theta_k, r_k) - g_0(\theta_k, r_k)) \right| \\
\leq \quad & \varepsilon^2 \left| \sum_{k=0}^{P_\beta - 1} \sum_{j=0}^{q-1} e^{-\lambda \varepsilon^2 (kq+j)} h(r_{kq+j})(g(\theta_{kq+j}, r_{kq+j}) - g_0(\theta_{kq+j}, r_{kq+j})) \right| \qquad (69) \\
+ \quad & \varepsilon^2 \left| \sum_{j=0}^{Q_\beta - 1} e^{-\lambda \varepsilon^2 (P_\beta q + j)} h(r_{P_\beta q+j})(g(\theta_{P_\beta q+j}, r_{P_\beta q+j}) - g_0(\theta_{P_\beta q+j}, r_{P_\beta q+j})) \right| .
\end{aligned}$$

On the one hand, we note that by (67) and $j \leq q < \varepsilon^{-b}$ we have

$$\begin{aligned}
\theta_{kq+j} &= \theta_{kq} + jr^* + \mathcal{O}(\varepsilon^{\beta - b}), \\
r_{kq+j} &= r^* + \mathcal{O}(\varepsilon^\beta).
\end{aligned}$$

Then for all $k \leq P_\beta$:

$$\begin{aligned}
& e^{-\lambda \varepsilon^2 (kq+j)} h(r_{kq+j})(g(\theta_{kq+j}, r_{kq+j}) - g_0(\theta_{kq+j}, r_{kq+j})) \\
= \quad & e^{-\lambda \varepsilon^2 kq} h(r_{kq})(g(\theta_{kq} + jr^*, r^*) - g_0(\theta_{kq} + jr^*, r^*)) + \mathcal{O}(e^{-\lambda \varepsilon^2 kq} \varepsilon^{1-b}).
\end{aligned}$$

Then:

$$\begin{aligned}
& \varepsilon^2 \left| \sum_{k=0}^{P_\beta - 1} \sum_{j=0}^{q-1} e^{-\lambda \varepsilon^2 (kq+j)} h(r_{kq+j})(g(\theta_{kq+j}, r_{kq+j}) - g_0(\theta_{kq+j}, r_{kq+j})) \right| \\
\leq \quad & \varepsilon^2 \left| \sum_{k=0}^{P_\beta - 1} e^{-\lambda \varepsilon^2 kq} h(r_{kN}) \sum_{j=0}^{q-1} (g(\theta_{kq} + jr^*, r^*) - g_0(\theta_{kq} + jr^*, r^*)) \right| \\
& + K \varepsilon^{3-2b} \sum_{k=0}^{P_\beta - 1} e^{-\lambda \varepsilon^2 kq}. \qquad (70)
\end{aligned}$$

Now, recalling that $r^* = p/q$, by the definition (68) of $g_0(\theta, r)$ for all $k < P_\beta$ we have:

$$\sum_{j=0}^{q-1} (g(\theta_{kq} + jr^*, r^*) - g_0(\theta_{kq} + jr^*, r^*)) = \sum_{j=0}^{q-1} g(\theta_{kq}, r^*) - q g_0(\theta_{kq}, r^*) = 0.$$

Moreover:

$$K \varepsilon^{3-2b} \sum_{k=0}^{P_\beta - 1} e^{-\lambda \varepsilon^2 kq} \leq K_\lambda \ \varepsilon^{1-2b}.$$

Using these estimates (70) yields:

$$\begin{gathered}\varepsilon^2 \left| \sum_{k=0}^{P_\beta - 1} \sum_{j=0}^{q-1} e^{-\lambda \varepsilon^2 (kq+j)} h(r_{kq+j})(g(\theta_{kq+j}, r_{kq+j}) - g_0(\theta_{kq+j}, r_{kq+j})) \right| \leq \\ K\,\varepsilon^{1-2b}.\end{gathered} \tag{71}$$

Note that $1 - 2b = 1 - \beta + \rho > 0$, since $\beta < 1$ and $\beta - 2b = \rho > 0$.

On the other hand, we have:

$$\begin{aligned}&\varepsilon^2 \left| \sum_{j=0}^{Q_\beta - 1} e^{-\lambda \varepsilon^2 (P_\beta N + j)} h(r_{P_\beta N+j})(g(_{P_\beta N+j}, r_{P_\beta N+j}) - g_0(r_{P_\beta N+j})) \right| \\ &\leq \varepsilon^2 K \sup_{(\theta,r)\in I_\beta} |h(r)(g(\theta,r) - g_0(\theta,r))| Q_\beta \leq \tilde{K}\varepsilon^{2-b}.\end{aligned} \tag{72}$$

Clearly, $2 - b > 0$. Substituting (71) and (72) in (69) yields item 1 of the Lemma.

Now let us consider item 2. If we take equation (70) and substitute $e^{-\lambda\varepsilon^2 k} h(r_k)$ by $h_k(r_k)$, we can write for all $k \leq P_\beta$:

$$\varepsilon^2 \left| \sum_{k=0}^{P_\beta - 1} \sum_{j=0}^{q-1} h_{kq+j}(r_{kq+j})(g(\theta_{kq+j}, r_{kq+j}) - g_0(\theta_{kq+j}, r_{kq+j})) \right| \tag{73}$$

$$\leq \quad \varepsilon^2 \left| \sum_{k=0}^{P_\beta - 1} h_{kq}(r_{kq}) \sum_{j=0}^{q-1} (g(\theta_{kq} + jr^*, r^*) - g_0(\theta_{kq} + jr^*, r^*)) \right| + K\varepsilon^{3-2b} P_\beta.$$

Again:

$$\sum_{j=0}^{q-1} (g(\theta_{kq} + jr^*, r^*) - g_0(\theta_{kq} + jr^*, r^*)) = 0,$$

and since $P_\beta < n_\beta < \varepsilon^{-2(1-\beta)-\delta}$ and $b = (\beta - \rho)/2$

$$K\,\varepsilon^{3-2b} P_\beta \leq K\,\varepsilon^{1+2\beta-2b} \leq K\varepsilon^{1+\rho+\beta}.$$

Then we have:

$$\begin{gathered}\varepsilon^2 \left| \sum_{k=0}^{P_\beta - 1} \sum_{j=0}^{q-1} e^{-\lambda \varepsilon^2 (kq+j)} h(r_{kq+j})(g(\theta_{kq+j}, r_{kq+j}) - g_0(\theta_{kq+j}, r_{kq+j})) \right| \leq \\ K\,\varepsilon^{1+\rho+\beta}.\end{gathered} \tag{74}$$

On the other hand we have:

$$\begin{aligned}&\varepsilon^2 \left| \sum_{j=0}^{Q_\beta - 1} h_{P_\beta N+j}(r_{P_\beta N+j})(g(_{P_\beta N+j}, r_{P_\beta N+j}) - g_0(r_{P_\beta N+j})) \right| \\ &\leq \varepsilon^2 K Q_\beta \leq \tilde{K}\varepsilon^{2-b}.\end{aligned} \tag{75}$$

We note that $2-b-2\beta>0$ for $b=(\beta-\rho)/2$ and $\beta<4/5$ and that $1-\beta>0$ if $\beta<1$. Thus, taking $\beta<4/5$ bounds (74) and (75) prove item 2 of the Lemma with $d=\min\{2-b-2\beta, 1-\beta, \rho\}>0$. □

Let r_0 belong to the IR case. Consider an interval $I_\beta=\{(\theta,r)\in\mathbb{T}\times\mathbb{R} : |r-r_0|\le\varepsilon^\beta\}$, for some $0<\beta<4/5$. Denote $n_\beta\in\mathbb{N}$ the exit time from I_β, that is the first number such that $(\theta_{n_\beta}, r_{n_\beta})\notin I_\beta$.

Lemma 5.5. *Let $f:\mathbb{R}\to\mathbb{R}$ be any $\mathcal{C}^l$ function with $l\ge 3$. Then, there exists $d>0$ such that for all $\lambda>0$ one has:*

$$\begin{aligned}&\mathbb{E}\Bigg(e^{-\lambda\varepsilon^2 n_\beta}f(r_{n_\beta})+\\ &\varepsilon^2\sum_{k=0}^{n_\beta-1}e^{-\lambda\varepsilon^2 k}\left[\lambda f(r_k)-\left(b(\theta_k,r_k)f'(r_k)+\frac{\sigma^2(\theta_k,r_k)}{2}f''(r_k)\right)\right]\Bigg)\\ &-f(r_0)=\mathcal{O}(\varepsilon^{2\beta+d}),\end{aligned}$$

where:

$$b(\theta,r)=\frac{1}{q}\sum_{i=0}^{q-1}E_2(\theta+ir,r),\qquad \sigma^2(\theta,r)=\frac{1}{q}\sum_{i=0}^{q-1}v^2(\theta+ir,r).$$

Proof. Let us fix any $0<\delta<1/6$. Again, denoting:

$$\begin{aligned}\eta=&\,e^{-\lambda\varepsilon^2 n_\beta}f(r_{n_\beta})+\\ &\varepsilon^2\sum_{k=0}^{n_\beta-1}e^{-\lambda\varepsilon^2 k}\quad\left[\lambda f(r_k)-\left(b(\theta_k,r_k)f'(r_k)+\frac{\sigma^2(\theta_k,r_k)}{2}f''(r_k)\right)\right]\end{aligned}\tag{76}$$

and using the law of total expectation, we have:

$$\begin{aligned}\mathbb{E}(\eta)=&\,\mathbb{E}\left(\eta\,|\,\varepsilon^{-(1-\beta)+\delta}\le n_\beta\le\varepsilon^{-2(1-\beta)-\delta}\right)\mathbb{P}\{\varepsilon^{-2(1-\beta)+\delta}\le n_\beta\le\varepsilon^{-2(1-\beta)-\delta}\}\\ &+\mathbb{E}\left(\eta\,|\,n_\beta<\varepsilon^{-(1-\beta)+\delta}\right)\mathbb{P}\{n_\beta<\varepsilon^{-2(1-\beta)+\delta}\}\\ &+\mathbb{E}\left(\eta\,|\,n_\beta>\varepsilon^{-2(1-\beta)-\delta}\right)\mathbb{P}\{n_\beta>\varepsilon^{-2(1-\beta)-\delta}\}.\end{aligned}\tag{77}$$

As in the proof of Lemma 5.3, for ε sufficiently small by Lemma B.2 and some $C>0$ independent of ε we have

$$\mathbb{P}\{n_\beta<\varepsilon^{-2(1-\beta)+\delta}\}\le\exp\left(-\frac{C}{\varepsilon^{2\delta}}\right),$$

and, thus, (77) yields:

$$\begin{aligned}\mathbb{E}(\eta)=&\,\mathbb{E}(\eta\,|\,\varepsilon^{-(1-\beta)+\delta}\le n_\beta\le\varepsilon^{-2(1-\beta)-\delta})\mathbb{P}\{\varepsilon^{-(1-\beta)+\delta}\le n_\beta\le\varepsilon^{-2(1-\beta)-\delta}\}\\ &+\mathbb{E}\left(\eta\,|\,n_\beta>\varepsilon^{-2(1-\beta)-\delta}\right)\mathbb{P}\{n_\beta>\varepsilon^{-2(1-\beta)-\delta}\}.\end{aligned}\tag{78}$$

Now, we can write:

$$\begin{aligned} e^{-\lambda\varepsilon^2 n_\beta} f(r_{n_\beta}) &= f(r_0) + \sum_{k=0}^{n_\beta-1} \Big[-\lambda\varepsilon^2 e^{-\lambda\varepsilon^2 k} f(r_k) + e^{-\lambda\varepsilon^2 k} f'(r_k)(r_{k+1}-r_k) \\ &\quad + \frac{1}{2} e^{-\lambda\varepsilon^2 k} f''(r_k)(r_{k+1}-r_k)^2 + \mathcal{O}(e^{-\lambda\varepsilon^2 k}\varepsilon^3) \Big], \end{aligned}$$

and then (76) can be rewritten as:

$$\begin{aligned} \eta = f(r_0) + & \\ \sum_{k=0}^{n_\beta-1} & \left[e^{-\lambda\varepsilon^2 k} f'(r_k)(r_{k+1}-r_k) + \frac{1}{2} e^{-\lambda\varepsilon^2 k} f''(r_k)(r_{k+1}-r_k)^2 \right] \\ -\varepsilon^2 \sum_{k=0}^{n_\beta-1} e^{-\lambda\varepsilon^2 k} & \left[b(\theta_k, r_k) f'(r_k) + \frac{\sigma^2(\theta_k, r_k)}{2} f''(r_k) \right] + \sum_{k=0}^{n_\beta-1} \mathcal{O}(e^{-\lambda\varepsilon^2 k}\varepsilon^3). \end{aligned} \tag{79}$$

Now, using (66) we have:

$$r_{k+1} - r_k = \varepsilon\omega_k[v(\theta_k, r_k) + \varepsilon v_2(\theta_k, r_k)] + \varepsilon^2 E_2(\theta_k, r_k) + \mathcal{O}(\varepsilon^{2+a}),$$

and:

$$(r_{k+1} - r_k)^2 = \varepsilon^2 v^2(\theta_k, r_k) + \mathcal{O}(\varepsilon^3).$$

Thus, (79) writes out as:

$$\begin{aligned} \eta &= f(r_0) + \sum_{k=0}^{n_\beta-1} e^{-\lambda\varepsilon^2 k} f'(r_k)\varepsilon\omega_k \left[v(\theta_k, r_k) + \varepsilon v_2(\theta_k, r_k) \right] \\ &\quad + \varepsilon^2 \sum_{k=0}^{n_\beta-1} e^{-\lambda\varepsilon^2 k} f'(r_k) \left[E_2(\theta_k, r_k) - b(\theta_k, r_k) \right] \\ &\quad + \frac{\varepsilon^2}{2} \sum_{k=0}^{n_\beta-1} e^{-\lambda\varepsilon^2 k} f''(r_k) \left[v^2(\theta_k, r_k) - \sigma^2(\theta_k, r_k) \right] \\ &\quad + \sum_{k=0}^{n_\beta-1} \mathcal{O}(e^{-\lambda\varepsilon^2 k}\varepsilon^{2+a}). \end{aligned} \tag{80}$$

Now we distinguish between the cases $\varepsilon^{-2(1-\beta)+\delta} \leq n_\beta \leq \varepsilon^{-2(1-\beta)-\delta}$ and $n_\beta > \varepsilon^{-2(1-\beta)-\delta}$. We shall start assuming that $\varepsilon^{-2(1-\beta)+\delta} \leq n_\beta \leq \varepsilon^{-2(1-\beta)-\delta}$. As in the proof of Lemma 5.3 we have:

$$\left| \sum_{k=0}^{n_\beta-1} \mathcal{O}(e^{-\lambda\varepsilon^2 k}\varepsilon^{2+a}) \right| \leq K\varepsilon^{2+a} n_\beta \leq K\varepsilon^{2\beta+a-\delta}. \tag{81}$$

We note that $1/6 - \delta > 0$ since we have taken $\delta < 1/6$. Now we use item 2 of Lemma 5.4 in (80) twice, taking first $h_k(r) = e^{-\lambda\varepsilon^2 k} f'(r)$ and $g(\theta, r) = E_2(\theta, r)$, and after $h_k(r) = e^{-\lambda\varepsilon^2 k} f''(r)$ and $g(\theta, r) = v^2(\theta, r)$. Then, using also (81), equation (80) yields:

$$\eta = f(r_0) + \sum_{k=0}^{n_\beta - 1} e^{-\lambda\varepsilon^2 k} f'(r_k)\varepsilon\omega_k \left[v(\theta_k, r_k) + \varepsilon v_2(\theta_k, r_k)\right] + \mathcal{O}(\varepsilon^{2\beta+d}), \qquad (82)$$

for some suitable $d > 0$.

Now turn to the case $n_\beta > \varepsilon^{-2(1-\beta)-\delta}$. The last term in (80) can be bounded by:

$$\left|\sum_{k=0}^{n_\beta - 1} \mathcal{O}(e^{-\lambda\varepsilon^2 k}\varepsilon^{2+a})\right| \leq K\varepsilon^{1+a} \sum_{k=0}^{n_\beta - 1} e^{-\lambda\varepsilon^2 k} = K\varepsilon^{1+a}\frac{1 - e^{-\lambda\varepsilon^2 n_\beta}}{1 - e^{-\lambda\varepsilon^2}} \leq K_\lambda \varepsilon^{a}, \qquad (83)$$

for some positive constants K and K_λ. Then, using item 1 of Lemma 5.4 twice (first with $h(r) = f'(r)$ and $g(\theta, r) = E_2(\theta, r)$, and later with $h(r) = f''(r)$ and $g(\theta, r) = v^2(\theta, r)$), we obtain the following bound for $n_\beta > \varepsilon^{-2(1-\beta)-\delta}$:

$$\eta = f(r_0) + \varepsilon \sum_{k=0}^{n_\beta - 1} e^{-\lambda\varepsilon^2 k} f'(r_k)\omega_k \left[v(\theta_k, r_k) + \varepsilon v_2(\theta_k, r_k)\right] + \mathcal{O}(\varepsilon^{d}). \qquad (84)$$

To finish the proof, we follow the same steps as in the proof of Lemma 5.3 and (78) yields:

$$\begin{aligned} & \mathbb{E}(\eta) - f(r_0) \\ = \; & \sum_{n\in\mathbb{N}} \mathbb{E}\left(\sum_{k=0}^{n-1} e^{-\lambda\varepsilon^2 k} f'(r_k)\varepsilon\omega_k \left[v(\theta_k, r_k) + \varepsilon v_2(\theta_k, r_k]\right)\right) \mathbb{P}\{n_\beta = n\} \\ + \; & \mathcal{O}(\varepsilon^{2\beta+d})\mathbb{P}\{\varepsilon^{-(1-\beta)+\delta} \leq n_\beta \leq \varepsilon^{-2(1-\beta)-\delta}\} + \mathcal{O}(\varepsilon^{d})\mathbb{P}\{n_\beta < \varepsilon^{-(1-\beta)+\delta}\} \\ = \; & \mathcal{O}(\varepsilon^{2\beta+d}), \end{aligned}$$

where by Lemma B.2 for some $C > 0$ independent of ε

$$\mathbb{P}\{n_\beta > \varepsilon^{-2(1-\beta)-\delta}\} = \exp\left(-\frac{C}{\varepsilon^{2\delta}}\right),$$

so that it is smaller than any power of ε. ☐

5.3 The Real Rational case

Here we study the system in the RR case, in the subdomain:

$$|r - p/q| \leq K_1 \varepsilon^{1/2}.$$



Figure 3: Level sets of the pendulum $p/q = 1/3$.

From the Normal Form Theorem, in the Real Rational strips the system takes the following form:

$$\begin{aligned} \theta_1 = &\quad \theta_0 + r_0 + \varepsilon\,[\mathbb{E}u(\theta_0, p/q) - \mathbb{E}v(\theta_0, p/q) + \mathbb{E}v_{p,q}(\theta_0, p/q) + E_3(\theta_0)] \\ + &\quad \varepsilon\omega_0 u(\theta_0, p/q) + \mathcal{O}(\varepsilon^{1+a}), \\ r_1 = &\quad r_0 + \varepsilon\mathbb{E}v_{p,q}(\theta_0, r_0) + \varepsilon\omega_0 v(\theta_0, r_0) \\ + &\quad \varepsilon^{3/2}\omega_0 v_2(\theta_0, r_0) + \varepsilon^{3/2}E_4(\theta_0, r_0) + \mathcal{O}(\varepsilon^{2+a}), \end{aligned} \tag{85}$$

where $v_2(\theta, r)$ can be written explicitly in terms of $v(\theta, r)$ and $S_1(\theta, r)$. The function E_4 is such that $\|E_4\|_{\mathcal{C}^0} \leq K$. We point out that it is a rescaled version of the function E_4 appearing in the Normal Form Theorem.

Recall also that:

$$\max\{\|\mathbb{E}u\|_{\mathcal{C}^0},\ \|\mathbb{E}v\|_{\mathcal{C}^0},\ \|\mathbb{E}v_{p,q}\|_{\mathcal{C}^0},\ \|E_3\|_{\mathcal{C}^0},\ \|u\|_{\mathcal{C}^0},\ \|v\|_{\mathcal{C}^0}\}\ \leq K.$$

Moreover, we have defined v_2 also in such a way that $\|v_2\|_{\mathcal{C}^0} \leq K$.

First, we switch to the resonant variable:

$$\hat{r} = r - p/q, \qquad 0 \leq |\hat{r}| \leq K_1\varepsilon^{1/2}.$$

With this new variable, system (85) writes out as:

$$\begin{aligned} \theta_1 = &\quad \theta_0 + p/q + \varepsilon\,[\mathbb{E}u(\theta_0, p/q) - \mathbb{E}v(\theta_0, p/q) + \mathbb{E}v_{p,q}(\theta_0, p/q) + E_3(\theta_0)] \\ + &\quad \hat{r}_0 + \varepsilon\omega_0 u(\theta_0, p/q) + \mathcal{O}(\varepsilon^{1+a}), \\ \hat{r}_1 = &\quad \hat{r}_0 + \varepsilon\hat{\mathbb{E}}v_{p,q}(\theta_0, \hat{r}_0) + \varepsilon^{3/2}\hat{E}_4(\theta_0, \hat{r}_0) + \varepsilon\omega_0\hat{v}(\theta_0, \hat{r}_0) \\ + &\quad \varepsilon^{3/2}\omega_0\hat{v}_2(\theta_0, \hat{r}_0) + \mathcal{O}(\varepsilon^{2+a}), \end{aligned} \tag{86}$$

where:

$$\begin{aligned} \hat{v}(\theta_0, \hat{r}_0) &= v(\theta_0, \hat{r}_0 + p/q), & \hat{v}_2(\theta_0, \hat{r}_0) &= v_2(\theta_0, \hat{r}_0 + p/q), \\ \hat{\mathbb{E}}v_{p,q}(\theta_0, \hat{r}_0) &= \mathbb{E}v_{p,q}(\theta_0, \hat{r}_0 + p/q), & \hat{E}_4(\theta_0, \hat{r}_0) &= E_4(\theta_0, \hat{r}_0 + p/q), \end{aligned}$$

From now on, we will abuse notation and drop all hats. We are interested in the $q-$th iteration of map (86), which is given by:

$$\begin{aligned}
\theta_q &= \theta_0 + qr_0 + \\
&+ \varepsilon \sum_{k=0}^{q-1} \left[\mathbb{E}u(\theta_k, p/q) - \mathbb{E}v(\theta_k, p/q) + (q-k)\mathbb{E}v_{p,q}(\theta_k, p/q) + E_3(\theta_k)\right] \\
&+ \varepsilon \textstyle\sum_{k=0}^{q-1} \omega_k \left[u(\theta_k, p/q) + v(\theta_k, p/q)\right] + \mathcal{O}(\varepsilon^{1+a}), \\
r_q &= r_0 + \varepsilon \sum_{k=0}^{q-1} \left[\mathbb{E}v_{p,q}(\theta_k, r_k) + \varepsilon^{1/2} E_4(\theta_k, r_k)\right] \\
&+ \varepsilon \textstyle\sum_{k=0}^{q-1} \omega_k \left[v(\theta_k, r_k) + \varepsilon^{1/2} v_2(\theta_k, r_k)\right] + \mathcal{O}(\varepsilon^{2+a}).
\end{aligned}$$

Taking into account that q is bounded for $0 \geq i \leq q$ we have:

$$\theta_i = \theta_0 + i(p/q + r_0) + \mathcal{O}(\varepsilon), \qquad r_i = r_0 + \mathcal{O}(\varepsilon),$$

we can rewrite the last system as:

$$\begin{aligned}
\theta_q =& \quad \theta_0 + qr_0 + \varepsilon \mathbb{E}u^{(q)}(\theta_0) + \varepsilon u^{(q)}(\theta_0, \omega_0^q) + \mathcal{O}(\varepsilon^{1+a}), \\
r_q =& \quad r_0 + \varepsilon \mathbb{E}v^{(q)}(\theta_0, r_0, \varepsilon) + \varepsilon v^{(q)}(\theta_0, r_0, \omega_0^q) + \varepsilon^{3/2} v_2^{(q)}(\theta_0, r_0, \omega_0^q) + \mathcal{O}(\varepsilon^2).
\end{aligned} \tag{87}$$

where $\omega_k^q = (\omega_{qk}, \ldots, \omega_{qk+q-1})$ and:

$$\begin{aligned}
\mathbb{E}u^{(q)}(\theta) &= \sum_{i=0}^{q-1} \left[\mathbb{E}u(\theta + ip/q, p/q) - \mathbb{E}v(\theta + ip/q, p/q) + \right. \\
&+ \left. (q-i)\mathbb{E}v_{p,q}(\theta + ip/q, p/q) + E_3(\theta + ip/q)\right], \\
u^{(q)}(\theta, \omega_k^q) &= \sum_{i=0}^{q-1} (q-i)\omega_{qk+i} \left[u(\theta + ip/q, p/q) + v(\theta + ip/q, p/q)\right], \\
\mathbb{E}v^{(q)}(\theta, r, \varepsilon) &= \sum_{i=0}^{q-1} \left[\mathbb{E}v_{p,q}(\theta + i(p/q + r), r) + \varepsilon^{1/2} E_4(\theta + i(p/q + r), r)\right], \\
v^{(q)}(\theta, r, \omega_k^q) &= \sum_{i=0}^{q-1} \omega_{qk+i} v(\theta + i(p/q + r), r), \\
v_2^{(q)}(\theta, r, \omega_k^q) &= \sum_{i=0}^{q-1} \omega_{qk+i} v_2(\theta + i(p/q + r), r).
\end{aligned}$$

Introduce a rescaled variable $r = R\sqrt{\varepsilon}$. Then (θ_1, R_1) are defined using system (87) with the corresponding rescaling. This can be rewritten in the following way, where we just keep the necessary ε-dependent terms:

$$\begin{aligned} \theta_q &= \theta_0 + qR_0\sqrt{\varepsilon} + \varepsilon \mathbb{E}u^{(q)}(\theta_0) + \varepsilon u^{(q)}(\theta_0, \omega_0^q) + \mathcal{O}(\varepsilon^{1+a}), \\ R_q &= R_0 + \varepsilon^{1/2}\mathbb{E}v^{(q)}(\theta_0, R_0\sqrt{\varepsilon}, 0) + \varepsilon^{1/2} v^{(q)}(\theta_0, R_0\sqrt{\varepsilon}, \omega_0^q) \\ &\quad + \varepsilon v_2^{(q)}(\theta_0, 0, \omega_0^q) + \mathcal{O}(\varepsilon^{3/2}). \end{aligned} \tag{88}$$

Consider the Hamiltonian:

$$H(\theta, R) = \frac{R^2}{2} - \frac{1}{q}\int_0^\theta \mathbb{E}v^{(q)}(s, R\sqrt{\varepsilon}, 0)ds. \tag{89}$$

Let $H_n := H(\theta_{qn}, R_{qn})$. We study the process: $(\theta_{qn}, H_n) := (\theta_n, H(\theta_{qn}, R_{qn})$, where (θ_{qn}, R_{qn}) is the process obtained iterating (88) n times. One can see that:

$$\begin{aligned} H_1 = H_0 + &\quad \sqrt{\varepsilon}R_0\, v^{(q)}(\theta_0, R_0\sqrt{\varepsilon}, \omega_0^q) \\ + &\quad \varepsilon F(\theta_0, R_0, \varepsilon) + \varepsilon G(\theta_0, R_0, \omega_0^q, \varepsilon) + \mathcal{O}(\varepsilon^{1+a}), \end{aligned} \tag{90}$$

where F and G are:

$$\begin{aligned} F(\theta, R) &= \frac{1}{q}\mathbb{E}v^{(q)}(\theta, 0, 0)\ \mathbb{E}u^{(q)}(\theta) \\ &- \frac{1}{q}\mathbb{E}v^{(q)}(\theta, 0, 0)\int_0^\theta \mathbb{E}v^{(q)}(s, 0, 0)ds \\ &+ \frac{q}{2}R^2\partial_\theta \mathbb{E}v^{(q)}(\theta, 0, 0) + \frac{1}{2}\left(\mathbb{E}v^{(q)}(\theta, 0, 0)\right)^2 \\ &+ \frac{1}{2}\sum_{i=0}^{q-1} v^2(\theta + ip/q, 0), \\ G(\theta, R, \omega_k^q) &= \frac{1}{q}\mathbb{E}v^{(q)}(\theta, 0, 0)\ u^{(q)}(\theta, \omega_k^q) \\ &- \frac{1}{q}v^{(q)}(\theta, 0, \omega_k^q, 0)\int_0^\theta \partial_r \mathbb{E}v^{(q)}(s, 0, 0)ds \\ +\frac{1}{2}\sum_{\substack{i,j=0 \\ i\neq j}}^{q-1} \omega_{qk+i} &\quad \omega_{qk+j} \quad v(\theta + ip/q, 0)v(\theta + jp/q, 0) \\ &+ \mathbb{E}v^{(q)}(\theta, 0, 0)v^{(q)}(\theta, 0, \omega_k^q) \\ &+ R\sum_{i=0}^{q-1}\omega_{qk+i}v_2(\theta + ip/q), 0). \end{aligned}$$

We note that since $|R| \leq K_1$ we have:

$$\|F\|_{\mathcal{C}^0} \leq K, \qquad \|G\|_{\mathcal{C}^0} \leq K.$$

Moreover, one has that for all $k \geq 0$:

$$\mathbb{E}(G(\theta_{qk}, R_{qk}, \omega_k^q)) = 0.$$

In terms of the variable H, the $\{|R| \leq K_1\}$ region can be written as:

$$I_{RR} := \{H \in \mathbb{R} \, : \, |H| \leq K_3\}, \tag{91}$$

for some constant K_3. Denote by n^* the stopping time of leaving I_{RR} for the first time.

Lemma 5.6. *Let $f : \mathbb{R} \to \mathbb{R}$ be any $\mathcal{C}^l$ function with $l \geq 3$. Then there exists $d > 2/5$ such that for all $\lambda > 0$ one has:*

$$\begin{aligned}&\mathbb{E}\Big(e^{-\lambda\varepsilon n^*} f(H_{n^*}) + \\ &\varepsilon \sum_{k=0}^{n^*-1} e^{-\lambda\varepsilon k}\left[\lambda f(H_k) - b(\theta_{qk}, R_{qk}) f'(H_k) - \frac{\sigma^2(\theta_{qk}, R_{qk})}{2} f''(H_k)\right]\Big) \\ &\qquad\qquad\qquad\qquad\qquad\qquad - f(H_0) = \mathcal{O}(\varepsilon^d),\end{aligned}$$

where:

$$b(\theta, R) = F(\theta, R), \quad \sigma^2(\theta, R) = R_{qk}^2 \sum_{i=0}^{q-1} v^2(\theta + ip/q, 0). \tag{92}$$

Proof. Depending on properties of the averaged potential $\mathbb{E}v^{(q)}(\theta, R\sqrt{\varepsilon}, 0)$ we decompose the region $\{|H| \leq K_1\}$ into several domains as follows.

In the interval $[-K_1, K_1]$ there are finitely many critical value of H due to the fact that $\mathbb{E}v^{(q)}$ is a trigonometric polynomial in θ. By assumption [**H5**] all critical points are nondegenerate. Therefore, there are finitely many of them. Consider a $\sqrt{\varepsilon}$-neighborhood of each one of them.

There are level sets that are not intersecting these neighbourhoods of the critical points. Call these level sets *regular.* The others levels intersect these neighbourhoods. Call these level sets *nearly critical.* Notice that the regular level sets are no necessarily connected (see Fig. 2).

Each connected component of a family of ovals gives rise to a segment and the union of such segments gives rise to a diffusion on a graph as in Freidlin-Wentzell [20], Sect. 8.

Now we study separately three regimes:

- a $\sqrt{\varepsilon}$-near a critical point;
- iterates with an initial condition on a near critical level set;
- iterates with an initial condition on a regular level set;

In the first case note that $n^* < \infty$ almost surely. Indeed, on the one hand if $(\theta^*, 0)$ is a critical point of the dominant part of the deterministic system, i.e. $\mathbb{E}u^{(q)}(\theta^*, 0, 0) = \mathbb{E}v^{(q)}(\theta^*, 0, 0) = 0$, then the drift of the process is given by:

$$b(\theta^*, 0) = \frac{1}{2}\sum_{i=0}^{q-1} v^2(\theta^* + ip/q, 0) > 0.$$

On the other hand, if $(\theta, 0)$ is not a critical point, then the drift of the R–component of the process (θ_{qn}, R_{qn}) defined in (88) is precisely, which is $\mathbb{E}v^{(q)}(\theta, 0, 0) \neq 0$. In conclusion, the process does not get stuck at $R = 0$. Since away from this zone the diffusion coefficient $\sigma(\theta, R)$ is nonzero, n^* must be finite with probability one.

In the second case denote:

$$\begin{aligned} \eta = {} & e^{-\lambda\varepsilon n^*} f(H_{n^*}) + \\ & \varepsilon \sum_{k=0}^{n^*-1} e^{-\lambda\varepsilon k}\left[\lambda f(H_k) - b(\theta_{qk}, R_{qk}) f'(H_k) - \frac{\sigma^2(\theta_{qk}, R_{qk})}{2} f''(H_k)\right]. \end{aligned} \tag{93}$$

Now we write:

$$\begin{aligned} e^{-\lambda\varepsilon n^*} f(H_{n^*}) &= f(H_0) + \sum_{k=0}^{n^*-1}\Big[-\lambda e^{-\lambda\varepsilon k} f(H_k) + e^{-\lambda\varepsilon k} f'(H_k)(H_{k+1} - H_k) \\ &\quad + \frac{1}{2} e^{-\lambda\varepsilon k} f''(H_k)(H_{k+1} - H_k)^2 + \mathcal{O}(e^{-\lambda\varepsilon k}\varepsilon^{3/2})\Big], \end{aligned}$$

so that (93) writes out as:

$$\begin{aligned} \eta = {} & f(H_0) + \\ & \sum_{k=0}^{n^*-1} \left[e^{-\lambda\varepsilon k} f'(H_k)(H_{k+1} - H_k) + \frac{1}{2} e^{-\lambda\varepsilon k} f''(H_k)(H_{k+1} - H_k)^2\right] \\ & -\varepsilon \sum_{k=0}^{n^*-1} e^{-\lambda\varepsilon k}\left[b(\theta_{qk}, R_{qk}) f'(H_k) + \frac{\sigma^2(\theta_{qk}, R_{qk})}{2} f''(H_k)\right] + \sum_{k=0}^{n^*-1} \mathcal{O}(e^{-\lambda\varepsilon k}\varepsilon^{3/2}). \end{aligned} \tag{94}$$

Now, using (90) it is clear that:

$$\begin{aligned} H_{k+1} - H_k = {} & \sqrt{\varepsilon} R_{qk} v^{(q)}(\theta_{qk}, R_{qk}\sqrt{\varepsilon}, \omega_k^q) + \\ & \varepsilon F(\theta_{qk}, R_{qk}) + \varepsilon G(\theta_{qk}, R_{qk}, \omega_k^q) + \mathcal{O}(\varepsilon^{1+a}). \end{aligned} \tag{95}$$

Moreover, we have:

$$\begin{aligned} (H_{k+1} - H_k)^2 &= \varepsilon R_{qk}^2 (v^{(q)}(\theta_{qk}, 0, \omega_k^q))^2 + \mathcal{O}(\varepsilon^{3/2}) \\ &= \varepsilon R_{qk}^2 \sum_{i=0}^{q-1} v^2(\theta_{qk} + ip/q, 0) + \varepsilon G_0(\theta_{qk}, R_{qk}, \omega_k^q) + \mathcal{O}(\varepsilon^{3/2}), \end{aligned} \tag{96}$$

where:

$$\begin{aligned}&G_0(\theta_{qk}, R_{qk}, \omega_k^q)\\ &= 2R_{qk}^2 \sum_{l=0}^{q-1} \sum_{j=l+1}^{q-1} \omega_{qk+l}\omega_{qk+j} v(\theta + lp/q, 0) v(\theta + jp/q, 0).\end{aligned}$$

We note that $\mathbb{E}(G_0(\theta_{qk}, R_{qk}, \omega_k^q)) = 0$.

Using (95) and (96), equation (94) writes out as:

$$\begin{aligned}\eta &= f(H_0) + \sqrt{\varepsilon} \sum_{k=0}^{n^*-1} e^{-\lambda\varepsilon k} f'(H_k) R_{qk} v^{(q)}(\theta_{qk}, \sqrt{\varepsilon} R_{qk}, \omega_k^q)\\ &\quad + \varepsilon \sum_{k=0}^{n^*-1} e^{-\lambda\varepsilon k} f'(H_k) \left[F(\theta_{qk}, R_{qk}) - b(\theta_{qk}, R_{qk})\right]\\ &\quad + \frac{\varepsilon}{2} \sum_{k=0}^{n^*-1} e^{-\lambda\varepsilon k} f''(H_k) \left[R_{qk}^2 \sum_{i=0}^{q-1} v^2(\theta_{qk} + ip/q, 0) - \sigma^2(\theta_{qk}, R_{qk})\right]\\ &\quad + \varepsilon \sum_{k=0}^{n^*-1} e^{-\lambda\varepsilon k} \left[f'(H_k) G(\theta_{qk}, R_{qk}, \omega_k^q) + \frac{f''(H_k)}{2} G_0(\theta_{qk}, R_{qk}, \omega_k^q)\right]\\ &\quad + \sum_{k=0}^{n^*-1} \mathcal{O}(e^{-\lambda\varepsilon k} \varepsilon^{1+a}).\end{aligned}$$

Now, on the one hand by definition of $b(\theta, R)$ and $\sigma^2(\theta, R)$ it is clear that:

$$\begin{aligned}F(\theta_{qk}, R_{qk}) - b(\theta_{qk}, R_{qk}) &= 0\\ R_{qk}^2 \sum_{i=0}^{q-1} v^2(\theta_{qk} + ip/q, 0) - \sigma^2(\theta_{qk}, R_{qk}) &= 0.\end{aligned} \tag{97}$$

On the other hand, the last term in (97) can be bounded by:

$$\begin{aligned}\left|\sum_{k=0}^{n^*-1} \mathcal{O}(e^{-\lambda\varepsilon k} \varepsilon^{1+a})\right| &\leq K\varepsilon^{1+a} \sum_{k=0}^{n^*-1} e^{-\lambda\varepsilon k} = K\varepsilon^{1+a} \frac{1 - e^{-\lambda\varepsilon n^*}}{1 - e^{-\lambda\varepsilon}}\\ &\leq K_\lambda \varepsilon^a,\end{aligned} \tag{98}$$

for some positive constants K and K_λ. Using (97), (97) and (98) in equation (97) we have:

$$\begin{aligned}\eta &= f(H_0) + \sqrt{\varepsilon} \sum_{k=0}^{n^*-1} e^{-\lambda\varepsilon k} f'(H_k) R_{qk} v^{(q)}(\theta_{qk}, \sqrt{\varepsilon} R_{qk}, \omega_k^q) \qquad (99)\\ &+ \varepsilon \sum_{k=0}^{n^*-1} e^{-\lambda\varepsilon k} \left[f'(H_k) G(\theta_{qk}, R_{qk}, \omega_k^q) + \frac{f''(H_k)}{2} G_0(\theta_{qk}, R_{qk}, \omega_k^q)\right] + \mathcal{O}(\varepsilon^a).\end{aligned}$$

Thus, to finish the proof, we just need to use that:

$$\mathbb{E}\left(v^{(q)}(\theta_{qk}, \sqrt{\varepsilon}R_{qk}, \omega_k^q)\right) = \mathbb{E}\left(G_0(\theta_{qk}, R_{qk}, \omega_k^q)\right) = \mathbb{E}\left(G(\theta_{qk}, R_{qk}, \omega_k^q)\right) = 0,$$

and that R_{qk} and H_k are independent of ω_k^q. Using these facts in (99), one obtains straightforwardly:

$$\mathbb{E}(\eta) - f(H_0) = \mathcal{O}(\varepsilon^a),$$

and the proof is finished. □

5.4 Transition Zones

Here we study the system in the RR case, in the subdomain:

$$K_1\varepsilon^{1/2} \le |r - p/q| \le K_2\varepsilon^{1/6},$$

for certain constants K_1 and K_2. From the Normal Form Theorem, in the Real Rational strips the system takes the following form:

$$\begin{array}{rcl} \theta_1 &=& \theta_0 + r_0 + \mathcal{O}(\varepsilon), \\ r_1 &=& r_0 + \varepsilon\mathbb{E}v_{p,q}(\theta_0, r_0) + \varepsilon\omega_0 v(\theta_0, r_0) + \mathcal{O}(\varepsilon^{3/2}). \end{array} \tag{100}$$

Recall that:

$$\|\mathbb{E}v_{p,q}\|_{\mathcal{C}^0} \le K, \qquad \|v\|_{\mathcal{C}^0} \le K.$$

For our purposes, it will be more convenient to work with the variable:

$$\hat{r} = r - p/q, \qquad K_1\varepsilon^{1/2} \le |\hat{r}| \le K_2\varepsilon^{1/6}.$$

With this new variable, system (100) writes out as:

$$\begin{array}{rcl} \theta_1 &=& \theta_0 + p/q + \hat{r}_0 + \mathcal{O}(\varepsilon), \\ \hat{r}_1 &=& \hat{r}_0 + \varepsilon\hat{\mathbb{E}}v_{p,q}(\theta_0, \hat{r}_0) + \varepsilon\omega_0 \hat{v}(\theta_0, \hat{r}_0) + \mathcal{O}(\varepsilon^{3/2}), \end{array} \tag{101}$$

where:

$$\hat{v}(\theta_0, \hat{r}_0) = v(\theta_0, \hat{r}_0 + p/q), \qquad \hat{\mathbb{E}}v_{p,q}(\theta_0, \hat{r}_0) = \mathbb{E}v_{p,q}(\theta_0, \hat{r}_0 + p/q).$$

From now on, we will abuse notation and drop all hats. We are interested in the $q-$th iteration of map (101), which is given by:

$$\begin{array}{rcl} \theta_q &=& \theta_0 + qr_0 + \mathcal{O}(\varepsilon), \\ r_q &=& r_0 + \varepsilon\displaystyle\sum_{k=0}^{q-1}\mathbb{E}v_{p,q}(\theta_k, r_k) + \varepsilon\sum_{k=0}^{q-1}\omega_k v(\theta_k, r_k) + \mathcal{O}(\varepsilon^{3/2}). \end{array}$$

Note that we have used that $|q|$ is bounded. Moreover, taking into account that for $i \leq q$, we have:

$$\theta_i = \theta_0 + i(p/q + r_0) + \mathcal{O}(\varepsilon), \qquad r_i = r_0 + \mathcal{O}(\varepsilon),$$

so that we can rewrite the last system as:

$$\begin{aligned} \theta_q &= \theta_0 + q r_0 + \mathcal{O}(\varepsilon), \\ r_q &= r_0 + \varepsilon \mathbb{E} v^{(q)}(\theta_0, r_0) + \varepsilon v^{(q)}(\theta_0, r_0, \omega_0^q) + \mathcal{O}(\varepsilon^{3/2}), \end{aligned} \tag{102}$$

where we use the notation $\omega_k^q = (\omega_{kq}, \ldots, \omega_{kq+q-1})$ and:

$$\begin{aligned} \mathbb{E} v^{(q)}(\theta, r) &= \sum_{i=0}^{q-1} \mathbb{E} v_{p,q}(\theta + i(p/q + r), r), \\ v^{(q)}(\theta, r, \omega_k^q) &= \sum_{i=0}^{q-1} \omega_{kq+i} v(\theta + i(p/q + r), r). \end{aligned}$$

We point out that for any $k \geq 0$:

$$\mathbb{E}\left(v^{(q)}(\theta_{qk}, r_{qk}, \omega_k^q)\right) = 0.$$

Now let us consider the following function:

$$H(\theta, r) = \frac{r^2}{2} - \frac{\varepsilon}{q} \int_0^\theta \mathbb{E} v^{(q)}(s, r) ds.$$

In this section we will study the process:

$$H_n := H(\theta_{qn}, r_{qn}),$$

where (θ_{qn}, r_{qn}) is the process obtained iterating (102) n times. One can easily see that:

$$H_1 = H_0 + \varepsilon r_0 v^{(q)}(\theta_0, r_0, \omega_0^q) + \mathcal{O}(\varepsilon r_0^2). \tag{103}$$

Now, given a constant $0 < \rho < 1/6$, we want to study the process H_n in the following $H-$strips:

$$I_\rho(r_0) = \{H \in \mathbb{R} : |H - H_0| \leq |r_0| \varepsilon^{1-\rho}\}. \tag{104}$$

We stress out that the width of these strips depends on the initial condition r_0 and ε. To avoid this, we can define the process:

$$\bar{H}_n := |r_0|^{-1} \varepsilon^{-1+\rho} H_n.$$

The process $\bar{H}_n$ is defined through:

$$\bar{H}_1 = \bar{H}_0 + \varepsilon^\rho v^{(q)}(\theta_0, r_0, \omega_0^q) + \mathcal{O}(\varepsilon^\rho |r_0|). \tag{105}$$

Clearly, if we denote by n_ρ the first exit time of the process H_n from the strip $I_\rho(r_0)$, it is also the first exit time of the process $\bar{H}_n$ of the strip:

$$I = \{H \in \mathbb{R} \,:\, |H - \bar{H}_0| \leq 1\}. \tag{106}$$

Lemma 5.7. *Let $f : \mathbb{R} \to \mathbb{R}$ be any $\mathcal{C}^l$ function with $l \geq 3$. Then there exists $d > 0$ such that for all $\lambda > 2/5$ one has:*

$$\mathbb{E}\left(e^{-\lambda\varepsilon^{2\rho} n_\rho} f(\bar{H}_{n_\rho}) + \varepsilon^{2\rho} \sum_{k=0}^{n_\rho - 1} e^{-\lambda\varepsilon^{2\rho}k}\left[\lambda f(\bar{H}_k) - \frac{\sigma^2(\theta_{qk}, r_{qk})}{2} f''(\bar{H}_k)\right]\right)$$
$$-f(\bar{H}_0) = \mathcal{O}(\varepsilon^d),$$

where:

$$\sigma^2(\theta, r) = \sum_{i=0}^{q-1} v^2(\theta + i(p/q + r), r).$$

Proof. Let us denote:

$$\eta = e^{-\lambda\varepsilon^{2\rho} n_\rho} f(r_{n_\rho}) + \varepsilon^{2\rho} \sum_{k=0}^{n_\rho - 1} e^{-\lambda\varepsilon^{2\rho}k}\left[\lambda f(\bar{H}_k) - \frac{\sigma^2(\theta_{qk}, r_{qk})}{2} f''(\bar{H}_k)\right]. \tag{107}$$

Now we write:

$$\begin{aligned} e^{-\lambda\varepsilon^{2\rho} n_\rho} f(\bar{H}_{n_\rho}) &= \\ f(\bar{H}_0) &+ \sum_{k=0}^{n_\rho - 1}\Big[-\lambda\varepsilon^{2\rho} e^{-\lambda\varepsilon^{2\rho}k} f(\bar{H}_k) + e^{-\lambda\varepsilon^{2\rho}k} f'(\bar{H}_k)(\bar{H}_{k+1} - \bar{H}_k) \\ &\quad + \frac{1}{2} e^{-\lambda\varepsilon^{2\rho}k} f''(\bar{H}_k)(\bar{H}_{k+1} - \bar{H}_k)^2 + \mathcal{O}(e^{-\lambda\varepsilon^2 k}\varepsilon^{3\rho})\Big], \end{aligned}$$

so that (107) writes out as:

$$\begin{aligned} \eta &= f(\bar{H}_0) + \\ \sum_{k=0}^{n_\rho - 1} &\quad \left[e^{-\lambda\varepsilon^{2\rho}k} f'(\bar{H}_k)(\bar{H}_{k+1} - \bar{H}_k) + \frac{1}{2} e^{-\lambda\varepsilon^{2\rho}k} f''(\bar{H}_k)(\bar{H}_{k+1} - \bar{H}_k)^2\right] \\ &\quad -\varepsilon^{2\rho} \sum_{k=0}^{n_\rho - 1} e^{-\lambda\varepsilon^{2\rho}k} \frac{\sigma^2(\theta_{qk}, r_{qk})}{2} f''(\bar{H}_k) + \sum_{k=0}^{n_\rho - 1} \mathcal{O}(e^{-\lambda\varepsilon^2 k}\varepsilon^{3\rho}). \end{aligned} \tag{108}$$

Now, using (105) it is clear that:

$$\bar{H}_{k+1} - \bar{H}_k = \varepsilon^\rho v^{(q)}(\theta_{qk}, r_{qk}, \omega_k^q) + \mathcal{O}(\varepsilon^\rho |r_{qk}|). \tag{109}$$

Moreover, we have:

$$\begin{aligned}(\bar{H}_{k+1}-\bar{H}_k)^2 &= \varepsilon^{2\rho}(v^{(q)}(\theta_{qk},r_{qk},\omega_k^q))^2+\mathcal{O}(\varepsilon^{2\rho}|r_{qk}|)\\ &= \varepsilon^{2\rho}\sum_{i=0}^{q-1}v^2(\theta_{qk}+i(p/q+r_{qk}),r_{qk})\\ &\quad +\varepsilon^{2\rho}G_0(\theta_{qk},r_{qk},\omega_k^q)+\mathcal{O}(\varepsilon^{2\rho}|r_{qk}|) \qquad (110)\end{aligned}$$

where:

$$G_0(\theta,r,\omega_k^q)=2\sum_{l=0}^{q-1}\sum_{j=l+1}^{q-1}\omega_{kq+l}\omega_{kq+j}v(\theta+l(p/q+r),r)v(\theta+j(p/q+r),r).$$

We note that $\mathbb{E}(G_0(\theta_{qk},r_{qk},\omega_k^q))=0$.

Using (109) and (110) and noting that $\varepsilon^{2\rho}|r_{qk}|\le\varepsilon^{\rho}|r_{qk}|$, equation (108) writes out as:

$$\begin{aligned}\eta &= f(\bar{H}_0)+\varepsilon^{\rho}\sum_{k=0}^{n_\rho-1}e^{-\lambda\varepsilon^{2\rho}k}f'(\bar{H}_k)v^{(q)}(\theta_{qk},r_{qk},\omega_k^q)\\ &\quad +\frac{\varepsilon^{2\rho}}{2}\sum_{k=0}^{n_\rho-1}e^{-\lambda\varepsilon^{2\rho}k}f''(\bar{H}_k)\left[\sum_{i=0}^{q-1}v^2(\theta_{qk}+i(p/q+r_{qk}),r_{qk})-\sigma^2(\theta_{qk},r_{qk})\right]\\ &\quad +\frac{\varepsilon^{2\rho}}{2}\sum_{k=0}^{n_\rho-1}e^{-\lambda\varepsilon^{2\rho}k}f''(\bar{H}_k)G_0(\theta_{qk},r_{qk},\omega_k^q)+\sum_{k=0}^{n_\rho-1}\mathcal{O}(e^{-\lambda\varepsilon^{2\rho}k}\varepsilon^{\rho}|r_{qk}|). \qquad (111)\end{aligned}$$

Now, on the one hand by definition of $\sigma^2(\theta,r)$ it is clear that:

$$\sum_{i=0}^{q-1}v^2(\theta_{qk}+i(p/q+r_{qk}),r_{qk})-\sigma^2(\theta_{qk},r_{qk})=0. \qquad (112)$$

On the other hand, the last term in (111) can be bounded by:

$$\begin{aligned}\left|\sum_{k=0}^{n_\rho-1}\mathcal{O}(e^{-\lambda\varepsilon^{2\rho}k}\varepsilon^{\rho}|r_{qk}|)\right| &\le K\varepsilon^{\rho+1/6}\sum_{k=0}^{n_\rho-1}e^{-\lambda\varepsilon^{2\rho}k}=K\varepsilon^{\rho+1/6}\frac{1-e^{-\lambda\varepsilon^{2\rho}n_\rho}}{1-e^{-\lambda\varepsilon^{2\rho}}}\\ &\le K_\lambda\varepsilon^{1/6-\rho}, \qquad (113)\end{aligned}$$

for some positive constants K and K_λ, where we have used that $|r_{qk}|\le\varepsilon^{1/6}$ if $0\le qk\le n_\rho$. Using (112) and (113) in equation (111), and denoting $d=1/6-\rho>0$, we have:

$$\begin{aligned}\eta &= f(\bar{H}_0)+\varepsilon^{\rho}\sum_{k=0}^{n_\rho-1}e^{-\lambda\varepsilon^{2\rho}k}f'(\bar{H}_k)v^{(q)}(\theta_{qk},r_{qk},\omega_k^q)\\ &\quad +\frac{\varepsilon^{2\rho}}{2}\sum_{k=0}^{n_\rho-1}e^{-\lambda\varepsilon^{2\rho}k}f''(\bar{H}_k)G_0(\theta_{qk},r_{qk},\omega_k^q)+\mathcal{O}(\varepsilon^d). \qquad (114)\end{aligned}$$

Thus, to finish the proof, we just need to use that:

$$\mathbb{E}\left(v^{(q)}(\theta_{qk}, r_{qk}, \omega_k^q)\right) = \mathbb{E}\left(G_0(\theta_{qk}, r_{qk}, \omega_k^q)\right) = 0.$$

Indeed, using this fact in (114), one obtains straightforwardly:

$$\mathbb{E}(\eta) - f(\bar{H}_0) = \mathcal{O}(\varepsilon^d),$$

and the proof is finished. □

A Measure of the domain covered by RR and IR intervals

In this section we show that, with the right choice of b, the measure of the the union of all strips of RR and IR type inside any compact set:

$$A_\beta = \cup_k I_\beta^k \subset \mathbb{T} \times B \qquad I_\beta^k \text{ strips of width } 2\varepsilon^\beta$$

goes to zero as $\varepsilon \to 0$.

In fact, we will do the proof for $A = [0,1]$. The general case is completely analogous. Let us consider:

$$\mathcal{R} = \{p/q \in \mathbb{Q} \,:\, p < q,\, \gcd(p,q) = 1,\, q < \varepsilon^{-b}\} = \cup_{q=1}^{q_{\max}} \mathcal{R}_q \subset [0,1],$$

where $q_{\max} = [\varepsilon^{-b}]$ and:

$$\mathcal{R}_q = \{p/q \in \mathbb{Q} \,:\, p < q,\, \gcd(p,q) = 1\}.$$

Finally we denote:

$$I_{\mathcal{R}} = \{I_\beta^k \subset [0,1] \,:\, \exists p/q \in \mathcal{R} \cap I_\beta\}.$$

Lemma A.1. *Let ρ be fixed, $0 < \rho < \beta$, and define $b = (\beta - \rho)/2$. Then, for each I_β such that there is at most one rational p/q satisfying $|q| \leq \varepsilon^{-b}$ the union $I_{\mathcal{R}}$ has the Lebesgue measure $\mu(I_{\mathcal{R}}) \leq \varepsilon^\rho$ and, therefore, as $\varepsilon \to 0$:*

$$\mu(I_{\mathcal{R}}) \to 0,$$

where μ denotes the Lebesgue measure.

Proof. On the one hand, suppose that $p/q \in I_\beta$, $q \leq \varepsilon^{-b}$. Then, for all $p'/q' \in I_\beta$, with p' and q' relatively prime and $p'/q' \neq p/q$, we have:

$$\varepsilon^\beta \geq |p/q - p'/q'| \geq \frac{1}{qq'} \geq \frac{\varepsilon^b}{q'}.$$

Therefore:

$$q' \geq \varepsilon^{-\beta+b} = \varepsilon^{-b-3\rho/2} > \varepsilon^{-b},$$

so the first part of the claim is proved.

On the other hand we note that, if $q_1 \neq q_2$, then $\mathcal{R}_{q_1} \cap \mathcal{R}_{q_2} = \emptyset$. Moreover, it is clear that $\#\mathcal{R}_q \leq q-1$ (and if q is prime then $\#\mathcal{R}_q = q-1$, so that the bound is optimal). Therefore we have:

$$\#\mathcal{R} \leq \sum_{q=1}^{q_{\max}} \#\mathcal{R}_q \leq \sum_{q=1}^{q_{\max}} q - 1 = \frac{q_{\max}^2}{2} < \varepsilon^{-2b}.$$

Since $\mu(I_\beta) = \varepsilon^\beta$, one has:

$$0 \leq \mu(I_{\mathcal{R}}) = \varepsilon^\beta \#\mathcal{R} < \varepsilon^\beta \varepsilon^{-2b} = \varepsilon^\rho,$$

so that the second claim of the lemma is also clear. □

B Sufficient condition for weak convergence and auxiliary lemmas

In order to prove that the r-component exhibits a diffusion process we need to adapt several lemmas from Ch. 8 sec. 3 [20]. We recall some terminology and notations (see Ch. 1 sec. 1 [20] for more details).

In the notations of section 1.6 we have

Lemma B.1. *(see Lm. 3.1, [20]) Let M be a metric space, Y a continuous mapping $M \mapsto Y(M)$, $Y(M)$ being a complete separable metric space. Let $(X_t^\varepsilon, P_x^\varepsilon)$ be a family of Markov processes in M; suppose that the process $Y(X_t^\varepsilon)$ has continuous trajectories. Let (y_t, P_y) be a Markov process with continuous paths in $Y(M)$ whose infinitesimal operator is A with domain of definition D_A. Suppose that the space $C[0,\infty)$ of continuous functions on $[0,\infty)$ with values in Γ is taken as the sample space, so that the distribution of the process in the space of continuous functions is simply P_y. Let Ψ be a subset of the space $C(Y(M))$ such that for measures μ_1, μ_2 on $Y(M)$ the equality $\int f d\mu_1 = \int f d\mu_2$ for all $f \in \Psi$ implies $\mu_1 = \mu_2$. Let D be a subset of D_A such that for every $f \in \Psi$ and $\lambda > 0$ the equation $\lambda F - AF = f$ has a solution $F \in D$.*

Suppose that for every $x \in M$ the family of distributions Q_x^ε of $Y(X_\bullet^\varepsilon)$ in the space $C[0,\infty)$ corresponding to the probabilities P_x^ε for all ε is tight; and that for every compact $K \subset Y(M)$, for every $f \in D$ and every $\lambda > 0$,

$$\mathbb{E}_x^\varepsilon \int_0^\infty \exp(-\lambda t) \left[\lambda f(Y(X_t^\varepsilon)) - Af(Y(X_t^\varepsilon))\right] dt \to f(Y(x))$$

as $\varepsilon \to 0$ uniformly in $x \in Y^{-1}(K)$.

Then Q_x^ε converges weakly as $\varepsilon \to 0$ to the probability measure $P_{Y(x)}$.

In our case $Y(M)$ is the real line. We use a discrete version of this lemma in our proof.

Similarly, to Lemma 3.2 [20] one can show that the family of distributions Q_x^ε (those of $Y(X_\bullet^\varepsilon)$ with respect to the probability measures P_x^ε in the space $C[0,\infty))$ with small nonzero ε is tight. Indeed, in our case speed of change of I is bounded. Denote $H(X)=H(r,\theta)=r^2/2$. Then

- for every $T>0$ and $\delta>0$ there exists H_0 such that
$$\mathbb{P}_x^\varepsilon\{\max_{0<t<T}|H(X_t^\varepsilon)|>H_0\}<\delta.$$
- for every compact subset $K\subset\mathbb{A}$ and for every sufficiently small $\rho>0$ there exists a constant A_ρ such that for every $a\in K$ there exists a function $f_\rho^a(y)$ on $Y(\mathbb{A})$ such that $f_\rho^a(a)\equiv 1, f_\rho^a(y)\equiv 0$ for $\rho(y,a)\geq\rho$, $0\leq f_\rho^a(y)\leq 1$ everywhere, and $f_\rho^a(Y(X_t^\varepsilon))+A_\rho t$ is a submartingale for all ε (see Stroock and Varadhan [37]).

In the proof we need an auxiliary lemmas. We study the random sums

$$S_n=\sum_{k=1}^n v_k\omega_k,\quad n\geq 1, \tag{115}$$

where $\{\omega_k\}_{k\geq 1}$ is a sequence of independent random variables with equal ± 1 with equal probability 1/2 each and $\{v_k\}_{k\geq 1}$ is a sequence such that

$$\lim_{n\to\infty}\frac{\sum_{k=1}^n v_k^2}{n}=\sigma.$$

Here is a standard

Lemma B.2. *$\{S_n/n^{1/2}\}_{n\geq 1}$ converges in distribution to the normal dirtribution $\mathcal{N}(0,\sigma^2)$.*

Recall that a characteristic function of a random variable X is a function $\phi_X:\mathbb{R}\to\mathbb{C}$ given by $\phi_X(t)=\mathbb{E}\exp(itX)$. Notice that it satisfies the following two properties:

- If X,Y are independent random variables, then $\varphi_{X+Y}=\varphi_X\cdot\varphi_Y$.
- $\varphi_{aX}(t)=\varphi_X(at)$.

A sufficient condition to prove convergence in distribution is as follows.

Theorem B.3 (Continuity theorem [6]). *Let $\{X_n\}_{n\geq 1},Y$ be random variables. If $\{\varphi_{X_n}(t)\}_{n\geq 1}$ converges to $\varphi_Y(t)$ for every $t\in\mathbb{R}$, then $\{X_n\}_{n\geq 1}$ converges in distribution to Y.*

A direct calculation shows that

$$\lim_{n\to\infty} \log \phi_{S_n/\sqrt{n}}(t) = -\frac{t^2}{2\sigma^2} \qquad \text{for all} \quad t \in \mathbb{R}.$$

This way of proof was communicated to the authors by Yuri Lima.

Acknowledgement: The authors warmly thank Leonid Koralov for numerious envigorating discussions of various topics involving stochatic processes. Communications with Dmitry Dolgopyat, Yuri Bakhtin, Jinxin Xue were useful for the project and gladly acknowledged by the authors. The second author acknowledges partial support of the NSF grant DMS-1402164.

References

[1] Arnold, V. I. Instabilities in dynamical systems with several degrees of freedom, Sov Math Dokl 5 (1964), 581–585;

[2] Arnold, V. I. Mathematical methods of classical mechanics, Graduate Texts in Mathematics, 60, Second Edition, Springer-Verlag, 1989.

[3] Arnold, V. I. Mathematical problems in classical physics. Trends and perspectives in applied mathematics, 1–20, Appl. Math. Sci., 100, Springer, New York, 1994.

[4] Bernard, P. The dynamics of pseudographs in convex Hamiltonian systems. *J. Amer. Math. Soc.*, 21(3):615–669, 2008.

[5] Bernard, P. Kaloshin, V. Zhang, K. Arnold diffusion in arbitrary degrees of freedom and 3-dimensional normally hyperbolic invariant cylinders, arXiv:1112.2773 [math.DS] 2011, 58pp, conditionally accepted to Acta Mathematica.

[6] Breiman, L. Probability Published by Society for Industrial and Applied Mathematics, 1992

[7] Brin, M. Stuck, G. Introduction to Dynamical Systems, Cambridge University Press, 2003.

[8] Cheng Ch.-Q., Arnold diffusion in nearly integrable Hamiltonian systems. arXiv: 1207.4016v2 9 Mar 2013, 127 pp;

[9] Cheng, Ch.-Q. Yan, J. Existence of diffusion orbits in a priori unstable Hamiltonian systems. *Journal of Differential Geometry*, 67 (2004), 457–517 & 82 (2009), 229–277;

[10] Chirikov. B. V. A universal instability of many-dimensional oscillator systems. Phys. Rep., 52(5): 264–379, 1979.

[11] Chirikov ,B.V. Vecheslavov, V.V. Theory of fast Arnold diffusion in many-frequency systems, J. Stat. Phys. 71(1/2): 243 (1993)

[12] de la Llave, R. Orbits of unbounded energy in perturbations of geodesic flows by periodic potentials. a simple construction preprint 70pp, 2005.

[13] Delshams, A. de la Llave, R. Seara, T. A geometric mechanism for diffusion in Hamiltonian systems overcoming the large gap problem: heuristics and rigorous verification on a model Mem. Amer. Math. Soc. 179 (2006), no. 844, pp.144

[14] de la Llave, Orbits of unbounded energy in perturbations of geodesic flows by periodic potentials. a simple construction preprint 70pp, 2005.

[15] Dolgopyat, D. Repulsion from resonance Memoires SMF, 128, 2012.

[16] Dumas, H. Laskar, J. Global Dynamics and Long-Time Stability in Hamiltonian via Numerical Frequency Analysis Phys Review Let. 70, no. 20, 1993, 2975–2979.

[17] Fejoz, J. Guardia, M. Kaloshin, V. Roldan, P. Kikrwood gaps and diffusion along mean motion resonance for the restricted planar three body problem, arXiv:1109.2892 [math.DS] 2013, to appear in Journal of the European Mathematical Society,

[18] Filonenko, N. Zaslavskii G. Stochastic instability of trapped particles and conditions of applicability of the quasi-linear approximation, Soviet Phys. JETP 27 (1968), 851–857.

[19] Freidlin, M. Sheu, S. Diffusion processes on graphs: stochastic differential equations, large deviation principle, Probability theory and related fields 116.2 (2000): 181–220;

[20] Freidlin, M. Wentzell, A. Random perturbations of dynamical systems, Grundlehren der Mathematischen Wissenschaften, Vol. 260, Springer, 2012, 258pp.

[21] Gidea, M. de la Llave. R Topological methods in the large gap problem. Discrete and Continuous Dynamical Systems, Vol. 14, 2006.

[22] Guardia, M. Kaloshin, V. Diffusive behavior for the generalized example of Arnold, in preparation

[23] Ibragimov, I. A. A note on the central limit theorems for dependent random variables, Theory of Probability and Its Applications, 1975.

[24] Kaloshin, V. Geometric proofs of Mather's accelerating and connecting theorems, Topics in Dynamics and Ergodic Theory (eds. S. Bezuglyi and S. Kolyada), London Mathematical Society, Lecture Notes Series, Cambridge University Press, 2003, 81—106

[25] Kaloshin, V. Zhang, K. A strong form of Arnold diffusion for two and a half degrees of freedom, arXiv:1212.1150 [math DS], 2012, 207pp,

[26] Kaloshin, V. Zhang, K. A strong form of Arnold diffusion for three and a half degrees of freedom, http://terpconnect.umd.edu/∼vkaloshi/ 36pp,

[27] Kaloshin, V. Zhang, K. Dynamics of the dominant Hamiltonians, with application to Arnold diffusion arXiv:1212.1150 [math DS], 2012, 83pp,

[28] Laskar, J. Frequency analysis for multi-dimensional systems. Global dynamics and diffusion, Physica D, 67 (1993), 257–281, North-Holland;

[29] Marco, J.-P. Sauzin, D. Wandering domains and random walks in Gevrey near integrable systems, Ergodic Theory & Dynamical Systems Vol. 24, 5 (volume spécial à la mémoire de M. Herman, édité par A. Fathi et J.-C. Yoccoz), 1619 – 1666, 2004.

[30] Moeckel, R. Transition tori in the five-body problem, J. Diff. Equations 129, 1996, 290–314.

[31] Moeckel, R. personal communications;

[32] Moser, J. Is the solar system stable? Math. Intelligencer, 1(2): 65–71, 1978/79.

[33] Piftankin G. Treshchev D. Separatrix maps in Hamiltonian systems, Russian Math. Surveys 62:2 219–322;

[34] Sauzin, D. Ergodicity and conservativity in the random iteration of standard maps, preprint 2006.

[35] Sauzin, D. Exemples de diffusion d'Arnold avec convergence vers un mouvement brownien, preprint 2006.

[36] Shatilov, D. Levichev, E. Simonov, E. and M. Zobov Application of frequency map analysis to beam-beam effects study in crab waist collision scheme Phys. Rev. ST Accel. Beams 14, January 2011

[37] Stroock, D.W. Varadhan, S.R.S. Multidimensional Diffusion Processes. Springer: Berlin, 1979.

[38] Treschev, D. Multidimensional Symplectic Separatrix Maps, J. Nonlinear Sciences 12 (2002), 27—58;

[39] Treschev, D. Evolution of slow variables in a priori unstable Hamiltonian systems Nonlinearity 17 (2004), no. 5, 1803–1841;

[40] Treschev, D Arnold diffusion far from strong resonances in multidimensional a priori unstable Hamiltonian systems Nonlinearity 25 (2012), no 9, 2717–2758.

[41] Wisdom, J. "The origin of the Kirkwood gaps - A mapping for asteroidal motion near the 3/1 commensurability". Astronomical Journal 87: 577–593, 1982.